

RESEARCH ARTICLE

Open Access



Recombinant truncated latency-associated peptide alleviates liver fibrosis in vitro and in vivo via inhibition of TGF- β /Smad pathway

Xudong Song^{1,2}, Jiayi Shi^{1,2}, Jieting Liu^{1,2}, Yong Liu^{1,3}, Yang Yu^{1,2}, Yufei Qiu^{1,2}, Zhiqin Cao^{1,2}, Yu Pan^{1,2}, Xiaohuan Yuan^{1,2}, Yanhui Chu^{1,2*} and Dan Wu^{1,2*} 

Abstract

Background: Liver fibrosis is a progressive liver injury response. Transforming growth factor β 1 (TGF- β 1) is over-secreted during liver fibrosis and promotes the development of liver fibrosis. Therapeutic approaches targeting TGF- β 1 and its downstream pathways are essential to inhibit liver fibrosis. The N-terminal latency-associated peptide (LAP) blocks the binding of TGF- β 1 to its receptor. Removal of LAP is critical for the activation of TGF- β 1. Therefore, inhibition of TGF- β 1 and its downstream pathways by LAP may be a potential approach to affect liver fibrosis.

Methods: Truncated LAP (tLAP) plasmids were constructed. Recombinant proteins were purified by Ni affinity chromatography. The effects of LAP and tLAP on liver fibrosis were investigated in TGF- β 1-induced HSC-T6 cells, AML12 cells and CCl₄-induced liver fibrosis mice by real time cellular analysis (RTCA), western blot, real-time quantitative PCR (RT-qPCR), immunofluorescence and pathological staining.

Results: LAP and tLAP could inhibit TGF- β 1-induced AML12 cells inflammation, apoptosis and EMT, and could inhibit TGF- β 1-induced HSC-T6 cells proliferation and fibrosis. LAP and tLAP could attenuate the pathological changes of liver fibrosis and inhibit the expression of fibrosis-related proteins and mRNAs in CCl₄-induced liver fibrosis mice.

Conclusion: LAP and tLAP could alleviate liver fibrosis in vitro and in vivo via inhibition of TGF- β /Smad pathway. TLAP has higher expression level and more effective anti-fibrosis activity compared to LAP. This study may provide new ideas for the treatment of liver fibrosis.

Keywords: Liver fibrosis, Transforming growth factor β 1, Latency-associated peptide, TGF- β /Smad pathway

Background

Liver fibrosis is a pathological change of chronic liver damage caused by various factors, such as alcohol consumption, viruses, and drug abuse, which can cause

chronic liver inflammation and lead to an abnormal trauma healing response (Bolognesi et al. 2007; Parola and Pinzani 2019; Su et al. 2014). The primary manifestation is the abnormal deposition of extracellular matrix (ECM) components, which ultimately leads to the formation of fibrous scarring (Chen et al. 2020; Theocharis et al. 2019). Liver fibrosis is a reversible pathological response, and timely reversal of liver fibrosis can prevent chronic liver disease from progressing to cirrhosis and liver cancer (Hernandez-Gea and Friedman 2011).

*Correspondence: chuyanhui@mdjmu.edu.cn; wudan@mdjmu.edu.cn

¹ Heilongjiang Province Key Laboratory of Anti-Fibrosis Biotherapy, Mudanjiang Medical University, No. 3, Tongxiang Street, Aimin District, Mudanjiang 157011, Heilongjiang, China
Full list of author information is available at the end of the article



The pathophysiological process of liver fibrosis is a complex process regulated by multiple signaling pathways, of which the TGF- β 1 signaling pathway plays a key role in the development of liver fibrosis (Lichtman et al. 2016; Xu et al. 2016). TGF- β 1 is lowly expressed in normal liver cells. However, when the liver is injured, TGF- β 1 is released to promote the activation of hepatic stellate cells (HSCs), which leads to the abnormal deposition of extracellular matrix, thus promoting the development of liver fibrosis (Ye et al. 2021; Zhang et al. 2016).

TGF- β 1 is a protein synthesized from an inactive precursor and requires activation to function (Khalil 1999). The TGF- β 1 proprotein consists of three parts: a signal peptide, an N-terminal precursor called a latency-associated peptide (LAP), and a C-terminal TGF- β 1. The signal peptide is cleaved before secretion and the proprotein becomes inactive pro-TGF- β 1 (Robertson et al. 2015). After cleavage by the Furin, the TGF- β homodimer binds with the LAP homodimer to form a small latent complex (SLC). SLC cross-links with latent TGF- β -binding protein (LTBP) to form the large latent complex (LLC). After the Arg-Gly-Asp (RGD) motif of LAP interacts with integrin α v β 6 or α v β 8, TGF- β 1 is released from latent complex and activates its receptor. Upon binding to TGF- β 1, TGF- β RII recruits and phosphorylates the intracellular domain of TGF- β RI, which subsequently activate downstream Smad signaling pathway (Shi et al. 2011).

LAP contains a straitjacket domain and an arm domain. The straitjacket is formed by the α 1 helix and the latent lasso. The arm domain contains two four-stranded β -sheets consisting of eight β -strands (Shi et al. 2011). The α 3 helix that connects the top and bottom strands is important for binding TGF- β 1 and forming latent complex (Stachowski et al. 2020). β -strand folds β 8 and β 9 connect two arm domains in a bowtie tail. The RGD motif in the arm domain is essential for integrin binding and TGF- β 1 activation (Dong et al. 2017). The α 2 helix and most of the β strands are conserved in all the TGF- β family members. The Binding of T β RI to TGF- β 1 is blocked by α 1, α 5 helix, fastener, β -strands β 1, β 3 and β 10. The binding of T β RII to TGF- β 1 is blocked by the latent lasso (Shi et al. 2011). The straitjacket and the arm domain together complete the wrapping of TGF- β 1. Removal of LAP is critical for TGF- β 1 activation (Saharinen et al. 1999). Inhibition of TGF- β 1 activation and receptor binding by LAP may be a potential approach to affect liver fibrosis.

In our previous study, the full-length PET28a-LAP recombinant plasmid was constructed and the LAP protein was purified by Ni affinity chromatography (Song et al. 2022). The soluble expression of LAP is low, which is not conducive to mass production and application. Therefore, based on the structural composition and

function of LAP, seven truncated LAP plasmids were constructed to screen truncated LAP (tLAP) with higher expression levels, lower molecular weight, and comparable activity to LAP. The effects of recombinant LAP and tLAP on liver fibrosis were investigated in vitro and vivo experiments.

Materials and methods

Plasmid construction and protein expression

LAP is composed of 249 amino acids (Human TGF- β 1 proprotein, P01137, 30-278). The truncated LAP fragments are No. 1 (1-132), No. 2 (1-65), No. 3 (1-100), No. 4 (45-132), No. 5 (77-249), No. 6 (77-193) and No.7 (194-249). Primers were designed and added with EcoRI and XhoI restriction sites. The target fragments were amplified by PCR and digested. The digested products were inserted into the PET-28a vector to construct recombinant plasmids with a His-tag at the N-terminus. The recombinant plasmids were transformed into *E. coli* C43(DE3) competent cells. Positive clones were cultured overnight at 37 °C in 3 mL LB medium. The bacterial liquid was inoculated into 20 mL of LB medium and cultured to an OD600 of 0.8. Then the bacterial liquid was added with 0.2 mM IPTG (Biofroxx, Germany) and incubated at 16 °C for 16 h. The collected cells were resuspended with 500 μ L Buffer A (25 mmol Tris 7.5, 300 mmol NaCl, 2 mM PMSF) and sonicated on ice. The supernatant was collected by centrifugation and combined with 30 μ L of Ni-NTA agarose at 4 °C for 1 h. 500 μ L Buffer B (25 mmol Tris 7.5, 300 mmol NaCl, 15 mmol Imidazole) was added to wash Ni-NTA agarose for 3 times. Protein expression was detected by 12% SDS-PAGE.

Protein purification and identification

The bacterial liquid cultured overnight was inoculated into 1 L LB medium at 1:200 and cultured at 37 °C for 4 h. Then the bacterial liquid was cooled to 16 °C and induced by 0.2 mM IPTG for 16 h. After sonication and centrifugation at 4 °C, the supernatant was loaded onto a Ni-NTA column. Impurity proteins were removed by washing buffer containing a gradient of imidazole. The target proteins were eluted with Buffer C (25 mmol Tris 7.5, 300 mmol NaCl, 300 mmol Imidazole). The imidazole was removed by multiple dialysis and the buffer was replaced with PBS. The proteins were detected by 12% SDS-PAGE and pulldown assay.

Cell culture

The hepatic stellate cell line, HSC-T6 (iCell Bioscience Inc, Shanghai, China) were cultured in DMEM medium (Cat.C11995500BT, Gibco, USA) supplemented with 1% penicillin/streptomycin (Cat. 15410, Gibco, BRL, USA) and

10% fetal bovine serum (FBS, Cat. P30-3302, PAN-Bio-tech, Germany). HSC-T6 cells were cultured at 37 °C in an incubator containing 5% CO₂.

The hepatocyte line, AML12 cells (iCell Bioscience Inc, Shanghai, China) were cultured in DME/F-12 medium (Cat.SH30023.01, HyClone, USA) supplemented with 1% penicillin/streptomycin and 10% fetal bovine serum. AML12 cells were cultured at 37 °C in an incubator containing 5% CO₂.

MTT and RTCA assay

The effects of recombinant LAP and tLAP on the proliferation of HSC-T6 cells induced by TGF-β1 were detected by MTT assay. Cells were plated in 96-well plates. HSC-T6 cells were treated with TGF-β1 (10 ng/mL, Pepro Tech, 100-21, USA), followed by the addition of 15, 30, and 60 μg/mL of LAP and tLAP, respectively. Absorbance was measured at OD490 after 24 h. Subsequently, the effects of recombinant proteins on the cell index of HSC-T6 cells were detected by RTCA. HSC-T6 cells were placed in E-Plate and TGF-β1 (10 ng/mL) was added. Recombinant LAP and tLAP were added, and the cell index was detected every 1 min for 72 h.

Immunofluorescence assay

About 50,000 cells were added to each well of the 6-well plate with cover glass by cell slide method, and TGF-β1 (10 ng/mL) was added to each well after the cells adhered. 60 μg/mL of LAP and tLAP were added for intervention. The cells were fixed with 4% paraformaldehyde after 48 h. The cells were treated with 0.1% TritonX-100 for 20 min and then sealed with 5% BSA at room temperature for 1 h. Cells were incubated overnight at 4 °C with the following primary antibodies from Affinity Biosciences (Cincinnati, OH, USA): α-SMA (Cat. AF1032), Collagen I (Cat. AF7001), FN (Cat.AF5335) at a dilution of 1:200. The next day, fluorescent secondary antibodies (Cat.A0568, Beyotime, China) were incubated at room temperature and protected from light. Nuclei were stained with DAPI. Cells were observed by confocal laser microscopy.

Animal experiments

Eight-week-old male C57BL/6 mice were purchased from Liaoning Changsheng Biotechnology Co., Ltd. (SCXK-2020-0001, China). The animal study protocol was approved by the Institutional Animal Care and Use Committee of Mudanjiang Medical University (IACUC-20210109-8). The mice were maintained at comfortable temperature on a 12 h:12 h light–dark cycle and 50–60% relative humidity. For the experiments, the mice were randomly divided into 4 groups. Control mice were injected with 1 mL/kg of olive oil twice a week for

8 weeks. The CCl₄ group was given 1 mL/kg of CCl₄ (prepared with olive oil, Cat. 56-23-5, Sigma-Aldrich, USA) twice a week for 8 weeks. The LAP and tLAP treatment group were administered CCl₄ in the same way as the CCl₄ group. At the same time, starting from week 7, mice were injected with LAP and tLAP proteins at a dose of 60 μg/mouse, twice a week for 2 weeks.

Histopathological analysis

The liver tissue was immediately fixed in a 10% formaldehyde solution and gradient ethanol for dehydration treatment, then embedded in paraffin to make 5 μm thick tissue sections. Before pathological staining, the slices were placed in an oven at 60 °C for 2 h, then dewaxed with xylene and rehydrated with gradient ethanol. Tissue sections were then stained with H&E, Sirius Red (Cat. MM1004, MaoKang biotechnology Co., Ltd, Shanghai, China), Masson (Solarbio, Peking, China), and were performed to observe the structural changes and collagen fiber deposition of the liver tissues in each group by an optical microscope.

For immunohistochemical staining, the tissue sections were dewaxed and rehydrated with gradient ethanol. After antigen repair with hydrogen peroxide and sodium citrate, the tissue sections were sealed with 5% BSA at room temperature for 1 h. Tissue sections were incubated overnight at 4 °C with the following primary antibodies from Affinity Biosciences (Cincinnati, OH, USA): α-SMA (Cat. AF1032), Collagen I (Cat. AF7001) at a dilution of 1:200. After washing 3 times by PBS, the anti-rabbit IgG (Cat. ZB-2301, Zhongshan-golden bridge, China) secondary antibodies was incubated at 37 °C for 1 h. DAB color solution was added for color development, and the nuclei were stained with hematoxylin. Tissue sections were observed with light microscopy and photographed.

Biochemical analysis

The blood samples were centrifuged at 3000 rpm at 4 °C for 15 min to separate the serum. The automatic biochemical analyzer was used to detect alanine aminotransferase (ALT) and aspartate aminotransferase (AST).

Reverse transcription quantitative PCR analysis

Total RNA was extracted from liver tissue and cells using HP Total RNA Kit (Cat. R6812-02, OMAGE, USA). Reverse transcription was performed using 1 μg total RNA via a first-strand cDNA synthesis kit (Cat.04897030001, Roche, Switzerland). StepOne real-time PCR system was used to detect gene expression levels in liver tissues and cells.

Western blot

Liver tissue or cells were lysed with RIPA lysate (Cat. R0020, Solarbio, China) at 4 °C for 30 min, and the supernatant was collected by centrifugation at 12,000 rpm/min at 4 °C for 10 min. Protein concentration was determined by BCA protein quantitation kit (Cat. 23225, Thermo, Germany). SDS-PAGE electrophoresis was performed with 50 µg protein per well, and the protein was transferred to PVDF membrane (Cat. IPVH00010, Millipore, USA). The PVDF membrane was sealed with 5% skim milk powder at room temperature for 1 h. Membranes were incubated overnight at 4°C. The following primary antibodies from Affinity Biosciences (Cincinnati, USA): α -SMA (Cat.AF1032), Collagen I (Cat.AF7001), FN (Cat. AF5335), β -actin (Cat. AF7018), GAPDH (Cat. T0004), IL-1 β (Cat.AF5103), IL-6 (Cat.DF6087), E-cadherin (Cat.AF0131), Bax (Cat.AF0120), Bcl-2 (Cat.AF6139); The following primary antibodies from Bioss Antibodies (Peking, China): Integrin Alpha V + β 6 (Cat.bs-5791R); The following primary antibodies from Cell Signaling Technology (Danvers, USA): Smad2 (Cat.3122), p-Smad2 (Cat.18338). After washing 3 times by TBST, the anti-rabbit IgG (Cat. ZB-2301, Zhongshan-golden bridge, China) and anti-mouse IgG (Cat. ZB-2305, Zhongshan-golden bridge, China) secondary antibodies was incubated at room temperature for 1 h. ECL chemiluminescence reagent (Cat. BL520A, Biosharp, China) was used for color rendering, and BIO-RAD gel imaging system and Image J was used for counting and statistical analysis of each band.

Statistical analysis

GraphPad Prism 8.0 software was used to analyze all the experimental data obtained in this study. The data are expressed as the mean \pm standard deviation. One-way ANOVA analyzed the differences among the groups. When $P < 0.05$, the data were considered statistically significant.

Results

Expression and purification of the recombinant LAP and tLAP

Seven truncated recombinant plasmids were constructed (Fig. 1A, B). The expression results showed that: No. 1 and No. 3 had certain protein expressions, but the expressions were lower than that of full-length LAP. The expression level of No.5 was much higher than that of full-length LAP (Fig. 1C). The structural region in No.5 is relatively conserved among members of the TGF- β family. The α 3 helix is associated with binding to TGF- β 1, and the α 5 helix, β 1, β 3 and β 10 are involved in wrapping

TGF- β 1 to block receptor binding. Therefore No. 5 was named as tLAP and selected for purification and functional experiments.


LAP and tLAP proteins were purified by Ni affinity chromatography. The results of SDS-PAGE showed that the expression level of tLAP was significantly higher than that of LAP. There are approximately 1.7 mg of LAP protein or 8 mg of tLAP protein in 1 L of bacterial solution (Fig. 2A, B). In previous studies, we demonstrated that recombinant LAP interacts with TGF- β 1 in vitro (Song et al. 2022). We explored whether recombinant tLAP interacts with TGF- β 1. The pull down results showed that TGF- β 1 could not bind to Ni-NTA agarose alone (lane 4). When TGF- β 1 was incubated with Ni-NTA agarose loaded with LAP and tLAP, TGF- β 1 could be detected in Ni-NTA agarose (lane 5, lane 6), suggesting that recombinant proteins may interact with TGF- β 1 (Fig. 2C).

Recombinant LAP and tLAP inhibited TGF- β 1-induced EMT, inflammation and apoptosis of AML12 cells

To explore the effects of recombinant LAP and tLAP on TGF- β 1-treated AML12 cells, we detected mRNA expression levels of apoptosis inflammation and EMT markers in TGF- β 1 (5 ng/mL)-treated AML12 cells. RT-qPCR results showed that the mRNA expression level of E-cadherin was decreased in AML12 cells under the action of TGF- β 1, while the expression level of α -SMA was significantly increased, these results suggest that AML12 cells lose the epithelial features and obtain mesenchymal phenotype by TGF- β 1. In addition, the mRNA expression of IL-1 β , IL-6, TNF- α and Bax in AML12 cells was significantly increased, while the expression of Bcl-2 was decreased. These results indicated that AML12 cells have inflammation and apoptosis under the action of TGF- β 1. When treated with recombinant LAP and tLAP, these effects caused by TGF- β 1 were inhibited (Fig. 3A). Western blot confirmed the expression of α -SMA, IL-1 β , IL-6 and Bax in AML12 cells was significantly increased, but the changes in EMT, inflammation and apoptosis markers after TGF- β 1 stimulation could inhibited by recombinant LAP and tLAP (Fig. 3B). Collectively, the results suggested that recombinant LAP and tLAP inhibited TGF- β 1-induced EMT, inflammation and apoptosis of AML12 cells, and recombinant tLAP showed more effective inhibition than LAP.

Recombinant LAP and tLAP inhibit TGF- β 1-induced HSC-T6 cell proliferation

HSC-T6 cells were treated with 10 ng/mL TGF- β 1 and different concentrations of LAP and tLAP. The MTT results showed that the proliferation of HSC-T6 cells significantly increased after TGF- β 1 stimulation compared with the control group. In contrast,



TGF-β1-induced HSC-T6 cell proliferation was significantly reduced after LAP and tLAP intervention (Fig. 4A). To confirm this result, the cell index (CI) of HSC-T6 cells was measured in real time every 1 min by using the RTCA system. The CI value of TGF-β1-induced cells was increased compared with the control

group, the CI value of cells induced by TGF-β1 was decreased after LAP and tLAP treatment (Fig. 4B). Since the CI value is positively correlated with cell viability. Therefore, MTT and RTCA results indicated that LAP and tLAP could inhibited TGF-β1-induced HSC-T6 cell proliferation, and tLAP showed more effective inhibitory activity than LAP.




Fig. 2 Purification and identification of Recombinant LAP and tLAP. **A** Purification of recombinant tLAP: lane M, protein marker; lane 1, Crushing bacterium fluid; lane 2, Supernatant; lane 3, Precipitation after crushing; lane 4, wash; lane 5–8, elution for 4 times. **B** Purification of recombinant LAP: lane M, protein marker; lane 1, Crushing bacterium fluid; lane 2, Supernatant; lane 3, Precipitation after crushing; lane 4, wash; lane 5–8, elution for 4 times. **C** Identification of recombinant tLAP by pull down assay. M, protein marker; lane 1, LAP + Ni-NTA agarose; lane 2, tLAP + Ni-NTA agarose; lane 3, TGF-β1; lane 4, TGF-β1 + Ni-NTA agarose; lane 5, TGF-β1 + LAP + Ni-NTA agarose. Lane 6, TGF-β1 + tLAP + Ni-NTA agarose

Recombinant LAP and tLAP inhibit TGF-β1-induced fibrosis in HSC-T6 cells

The effects of LAP and tLAP on the expression of fibrosis-related mRNA in TGF-β1-induced HSC-T6 cells were detected by RT-qPCR. The RT-qPCR results showed that the mRNA expression of α-SMA, Collagen I, Collagen IV, and Fibronectin in the TGF-β1 group were significantly increased compared with the control group, compared with the TGF-β1 group, the mRNA expression were significantly decreased by LAP and tLAP (Fig. 4C). The western blot results showed that compared with the control group, the expression of α-SMA, Collagen I and Fibronectin in HSC-T6 were significantly increased by TGF-β1 stimulation. Compared with the TGF-β1 group, the expression of fibrosis-related proteins were significantly reduced in LAP and tLAP groups (Fig. 4D). The immunofluorescence results showed that after TGF-β1 treatment, the fluorescence intensity of α-SMA, Collagen I and Fibronectin was significantly enhanced. The fibrosis-related protein fluorescence intensity decreased after LAP and tLAP treatment (Fig. 5A–C). The results showed recombinant LAP and tLAP can inhibit TGF-β1-induced fibrosis in HSC-T6 cells, and compared with recombinant LAP, tLAP has more effective anti-fibrotic activity.

Recombinant LAP and tLAP alleviate CCl4-induced liver fibrosis in mice

The anti-fibrotic activity of LAP and tLAP in vivo was observed by CCl4-induced liver fibrosis model in C57BL/6 mice. The mRNA expression of IL-1β,

IL-6, TNF-α, TGF-β, α-SMA, Collagen I, Collagen IV, and Fibronectin in the CCl4 group were significantly increased compared with the control group, demonstrated liver inflammation and fibrosis. The recombinant LAP and tLAP could effectively inhibit the above gene changes induced by CCl4, and recombinant tLAP had more significant inhibition than LAP (Fig. 6A). Western blot confirmed that recombinant LAP and tLAP inhibited liver inflammation and fibrosis associated marker proteins elevation induced by CCl4 (Fig. 6B).

The results of H&E staining showed that the hepatocytes in the CCl4 model group had obvious inflammatory infiltration, deformation, and necrosis, which were alleviated in LAP and tLAP groups. Masson staining and Sirius red staining showed that compared with the control group, the liver of the CCl4 model group had a large number of collagen fiber hyperplasia, while the LAP and tLAP groups significantly decreased the fibrous hyperplasia. The expression of α-SMA and Collagen I in liver was analyzed by immunohistochemical staining. The results showed that compared with the model group, the expression of α-SMA and Collagen I were significantly decreased in the LAP and tLAP groups (Fig. 7A, B). Biochemical analysis showed that the serum ALT and AST of mice in the CCl4 group were significantly increased than those in the control group. Compared with the model group, serum ALT and AST were significantly decreased by LAP and tLAP (Fig. 7C). These results suggested that recombinant LAP and tLAP can effectively alleviate CCl4-induced liver fibrosis in C57BL/6 mice.





Fig. 3 Detection of the effects of LAP and tLAP on TGF-β1-induced apoptosis and EMT in AML12 cells. **A** Detection of the mRNA expression of Bax, Bcl-2, IL-6, IL-1β, TNF-α, TGF-β, α-SMA and E-cadherin in AML12 cells. **B** Detection of the expression of Bax, Bcl-2, IL-6, IL-1β, α-SMA and E-cadherin in AML12 cells. ***P < 0.001 vs control group; #P < 0.05 vs TGF-β1 group; ##P < 0.01 vs TGF-β1 group; ###P < 0.001 vs TGF-β1 group; ΔP < 0.05, TGF-β1 + tLAP group vs TGF-β1 + LAP group; ΔΔP < 0.01, TGF-β1 + tLAP group vs TGF-β1 + LAP group; ΔΔΔP < 0.001, TGF-β1 + tLAP group vs TGF-β1 + LAP group

(See figure on next page.)

Fig. 4 Detection of the effect of LAP and tLAP on liver fibrosis in HSC-T6 cells. **A** Detection of the effect of LAP and tLAP on cell proliferation by MTT. **B** Detection of the effect of LAP and tLAP on cell index by RTCA. **C** Detection of the mRNA expression α-SMA, Collagen I, Collagen IV, and Fibronectin in HSC-T6 cells. **D** Detection of the expression of α-SMA, Collagen I and Fibronectin in HSC-T6 cells. **P < 0.01 vs control group; ***P < 0.001 vs control group; #P < 0.05 vs TGF-β1 group; ##P < 0.01 vs TGF-β1 group; ###P < 0.001 vs TGF-β1 group; ΔP < 0.05, TGF-β1 + tLAP group vs TGF-β1 + LAP group; ΔΔP < 0.01, TGF-β1 + tLAP group vs TGF-β1 + LAP group



Recombinant LAP and tLAP alleviate liver fibrosis via inhibition of TGF- β /Smad pathway

These results above indicated that LAP and tLAP could alleviate liver fibrosis in vitro and in vivo. However, liver fibrosis is a complex process regulated by multiple signaling pathways. LAP interacts with a variety of proteins in vivo. How to determine whether its anti-fibrosis activity is related to TGF- β 1? Previous studies have found that LAP contains an arginine-glycine-aspartic acid (RGD) motif, making LAP a ligand for integrin $\alpha\beta$ 6. Integrins bind to LAP through the RGD motif to regulate the activation of TGF- β 1 (Thomas et al. 2002). Therefore, we examined the expression of $\alpha\beta$ 6 in HSC-T6 cells and liver, and the results showed that the mRNA expression of $\alpha\beta$ 6 was significantly increased in fibrotic HSC-T6 cells and liver, while recombinant LAP and tLAP inhibited the expression of $\alpha\beta$ 6 (Fig. 8A–C). In addition, the effects of LAP and tLAP on the TGF- β 1 downstream Smad pathway were investigated in HSC-T6 cells and liver. The results showed that the expression level of p-Smad2 protein was significantly increased in fibrotic HSC-T6 cells and liver, and recombinant LAP and tLAP could effectively inhibit the phosphorylation of smad2 protein. (Fig. 8D, E). These results suggested that recombinant LAP and tLAP may alleviate liver fibrosis via inhibition of TGF- β /Smad pathway.

Discussion

When the liver suffers sustained injury, it often induces a self-repair response, leading to fibrous connective tissue deposition and eventually liver fibrosis. If not treated in time, most patients will progress to liver cirrhosis, liver failure, or even liver cancer (Ramadori and Saile 2004; Tanwar et al. 2020). Active and effective intervention is expected to delay or prevent the progression of liver fibrosis, prolong the survival of patients, and improve the quality of life of patients (Campana and Iredale 2017).

TGF- β 1 is a highly pleiotropic cytokine with a wide range of biological functions. TGF- β 1 signal transduction is involved in almost all stages of liver disease progression, including inflammation, fibrosis, cirrhosis, and even liver cancer (Fabregat et al. 2016). TGF- β 1 promotes the activation of hepatic stellate cells into myofibroblasts and stimulates ECM synthesis (Shrestha et al. 2016). Studies have shown that serum levels of TGF- β 1 are significantly higher in patients with liver fibrosis than in healthy people (Cayon et al. 2006). TGF- β 1 is often secreted in the form of a latent complex. Most TGF- β 1 is synthesized by

parenchymal cells and secreted to the extracellular space, and the whole process is affected by a variety of factors. The signal peptide is cleaved before secretion. Under the action of enzyme, a complex composed of an N-terminal LAP and C-terminal mature TGF- β is connected by a non-covalent bond. Then, the complex becomes extracellular by changing pH, ion strength, or protease hydrolysis. Stripping LAP or changing LAP conformation releases mature TGF- β . When TGF- β binds to LAP, it cannot bind to its receptor (Clark and Coker 1998). The extracellular matrix glycoprotein thrombus reactive protein-1 (TSP-1) is one of the main physiological activators of TGF- β (Murphy-Ullrich and Poczatek 2000). TSP-1 activates TGF- β 1 by binding to LAP, thereby disrupting the domain binding of LAP-mature TGF- β 1 (Ribeiro et al. 1999). Thus, LAP removal is an essential step in TGF- β 1 activation. Affecting TGF- β 1 activation and receptor binding by LAP may be a potential treatment for liver fibrosis.

Considering the importance of LAP in TGF- β 1 activation, LAP may be a key target for inhibiting the TGF- β 1 pathway. In this study, we examined the anti-fibrotic effects of recombinant LAP and tLAP in vitro and in vivo. The anti-fibrosis activity of recombinant LAP and tLAP was confirmed based on the following results: (1) Recombinant LAP and tLAP inhibited TGF- β 1-induced EMT, inflammation and apoptosis of AML12 cells; (2) Recombinant LAP and tLAP inhibit TGF- β 1-induced proliferation and fibrosis of HSC-T6 cells; (3) Recombinant LAP and tLAP have protective effect on CCl₄-induced liver injury and liver fibrosis; (4) Recombinant LAP and tLAP inhibit the phosphorylation of Smad2 protein.

In our previous study, the LAP-pET28a expression vector was constructed, and the recombinant protein was successfully expressed and purified. But the soluble expression of LAP was low, which was not conducive to large-scale production and application. Therefore, in order to improve its yield, based on the structural composition and function of LAP, a truncated tLAP-pET28a plasmid was constructed. By screening the induction conditions, we successfully purified the recombinant tLAP. The results showed that the yield of recombinant tLAP is nearly 5 times that of LAP.

We investigated the effects of recombinant LAP and tLAP on TGF- β 1-induced AML12 cells. Our results showed expression of α -SMA, IL-1 β , IL-6, TNF- α and Bax in AML12 cells treated with TGF- β 1 were significantly increased and expression of E-cadherin and Bcl-2

(See figure on next page.)

Fig. 5 Detection of fibrosis-related proteins expression in HSC-T6 cells by Immunofluorescence. **A** Immunofluorescence detection of α -SMA. **B** Immunofluorescence detection of collagen I. **C** Immunofluorescence detection of Fibronectin. DAPI (blue, nuclear stain) and antibodies to α -SMA, Collagen I and Fibronectin (red), immunofluorescence staining (magnification \times 200), Scale bars = 40 μ m. **P < 0.01 vs control group; ***P < 0.001 vs control group; #P < 0.05 vs TGF- β 1 group; ##P < 0.01 vs TGF- β 1 group





Fig. 5 (See legend on previous page.)



were decreased, confirmed the occurrence of EMT, inflammation and apoptosis. Recombinant LAP and tLAP significantly improved TGF-β1-induced EMT, inflammation and apoptosis in AML12 cells. Then, we detected the effects of recombinant LAP and tLAP on the

proliferation of HSC-T6 cells induced by TGF-β1 using MTT and RTCA. The results showed that recombinant LAP and tLAP can significantly reduce TGF-β1-induced HSC-T6 cell proliferation. Previous studies have shown that TGF-β1 treatment can activate the TGF-β/Smad





Fig. 7 H&E, Sirius Red, Masson's Trichrome Staining and Immunohistochemical and Liver Function Indicators. **A** H&E, Sirius Red, Masson staining, and immunohistochemistry of the liver tissues. **B** Detection of serum ALT and AST activity in C57BL/6 mice. ***P < 0.001 vs control group; #P < 0.05 vs CCl₄ group; ##P < 0.01 vs CCl₄ group. ###P < 0.001 vs CCl₄ group

pathway in HSC-T6 cells, leading to overexpression of fibrosis related genes and proteins (Gong et al. 2020). In this study, HSC-T6 cells were stimulated with TGF-β1 and treated with recombinant LAP and tLAP. The RT-qPCR, western blot and immunofluorescence results

showed that recombinant LAP and tLAP could reduce the expression of fibrosis related genes and proteins in HSC-T6 cells induced by TGF-β1, and the results of RT-qPCR showed that tLAP had more effective anti-fibrosis activity compared to LAP.



In order to study the anti-fibrosis activity of recombinant LAP and tLAP in vivo. The liver fibrosis model was established by CCl₄. It has been confirmed that CCl₄ can induce liver inflammation, destroy liver cell membrane, cause liver lobule cell necrosis, and lead to liver fibrosis in mice (Unsal et al. 2021). AST and ALT are common markers of liver injury. Experimental results showed that recombinant LAP and tLAP could reduce the expression of ALT and AST in hepatic fibrosis mice. In addition, HE staining showed that recombinant LAP and tLAP could significantly reduce the infiltration and aggregation of liver inflammatory cells caused by CCl₄, and MASSON and Sirius red staining showed that recombinant LAP

and tLAP could significantly reduce the area of collagen fibers around the portal area of fibrosis liver. RT-qPCR and western blot analysis showed that recombinant LAP and tLAP could significantly inhibit the increased expression of fibrosis related genes and proteins in liver induced by CCl₄, and consistent with the results of in vitro experiments, the recombinant tLAP protein showed more effective anti-fibrosis activity on the gene expression level.

Previous studies have shown that $\alpha v\beta 6$ is related to the activation of TGF- $\beta 1$, and LAP interacts with $\alpha v\beta 6$ (Thomas et al. 2002). Therefore, the expression of $\alpha v\beta 6$ was detected in this study. The results showed that

recombinant LAP and tLAP can inhibit the expression of $\alpha\beta6$ in fibrotic cells and liver. When TGF- β 1 binds to HSCs membrane receptors and activates TGF β RI, the activated TGF β RI can induce Smad2 and Smad3 phosphorylation and combine with Smad4 to form smad2/3/4 complex, then transported into the nucleus together and regulates gene transcription (Dewidar et al. 2019). To determine whether the effects of LAP and tLAP on liver fibrosis are related to TGF- β /Smad pathways. We investigated the expression levels of Smad2 and p-Smad2 in HSC-T6 cells and liver. Results showed that p-Smad2 protein expression levels were significantly increased in fibrotic livers and cells, and recombinant LAP and tLAP inhibited Smad2 protein phosphorylation. These results suggested that recombinant LAP and tLAP may alleviate liver fibrosis through the TGF- β /Smad signaling pathway.

Conclusions

In this study, we successfully constructed and prepared recombinant tLAP. The results indicated that recombinant LAP and tLAP could alleviate liver fibrosis via inhibition of TGF- β /Smad pathway. TLAP has higher expression level and more effective anti-fibrosis activity compared to LAP. This study may provide new ideas for the treatment of liver fibrosis.

Abbreviations

TGF- β 1: Transforming growth factor- β 1; LAP: Latency-associated peptide; HSCs: Hepatic stellate cells; RTCA: Real time cellular analysis; RT-qPCR: Real-time quantitative PCR; ECM: Extracellular matrix; SLC: Small latent complex; LTBP: Latent TGF- β -binding protein; LLC: Large latent complex; tLAP: Truncated LAP; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase.

Acknowledgements

We thank the Department of Laboratory Animals in Mudanjiang Medical University for supporting the animal work.

Author contributions

Methodology, DW and YHC; software, XDS, JYS; formal analysis, XHY; investigation, XDS, YY, YFQ, ZQC and YP; resources, DW, YHC and YL; data curation, DW and XDS; writing—original draft preparation, DW and XDS; writing—review and editing, DW; visualization, DW; supervision, YHC and XHY; project administration, XDS, JYS; funding acquisition, DW, YHC and JTL. All authors read and approved the final manuscript.

Funding

This study was funded by Central Finance supports Local Colleges, and Universities Talent Development Funding from Heilongjiang Provincial Department of Finance (High Level Talent Support Project) (Grant No. 2020GSP09); Natural Science Foundation of Heilongjiang Province (Grant No. YQ2020H026); the Basic Scientific Research Project of University belongs to Heilongjiang (Grant No. 2018-KYYWF-0136, Grant No. 2018-KYYWFM-0044); National Natural Science Foundation of China (Grant No. 81770828).

Availability of data and materials

The data used and analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The animal study protocol was approved by the Institutional Animal Care and Use Committee of Mudanjiang Medical University (IACUC-20210109-8).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Heilongjiang Province Key Laboratory of Anti-Fibrosis Biotherapy, Mudanjiang Medical University, No. 3, Tongxiang Street, Aimin District, Mudanjiang 157011, Heilongjiang, China. ²College of Life Sciences, Mudanjiang Medical University, Mudanjiang 157011, Heilongjiang, China. ³Mudanjiang Medical University, Mudanjiang 157011, Heilongjiang, China.

Received: 6 March 2022 Accepted: 7 July 2022

Published online: 16 July 2022

References

- Bolognesi M, Sacerdoti D, Mescoli C, Bombonato G, Cillo U, Merenda R, Giacomelli L, Merkel C, Ruge M, Gatta A. Different hemodynamic patterns of alcoholic and viral endstage cirrhosis: analysis of explanted liver weight, degree of fibrosis and splanchnic Doppler parameters. *Scand J Gastroenterol.* 2007;42:256–62.
- Campana L, Iredale JP. Regression of liver fibrosis. *Semin Liver Dis.* 2017;37:1–10.
- Cayon A, Crespo J, Mayorga M, Guerra A, Pons-Romero F. Increased expression of Ob-Rb and its relationship with the overexpression of TGF-beta1 and the stage of fibrosis in patients with nonalcoholic steatohepatitis. *Liver Int.* 2006;26:1065–71.
- Chen Y, Fan Y, Guo DY, Xu B, Shi XY, Li JT, Duan LF. Study on the relationship between hepatic fibrosis and epithelial-mesenchymal transition in intrahepatic cells. *Biomed Pharmacother.* 2020;129: 110413.
- Clark DA, Coker R. Transforming growth factor-beta (TGF-beta). *Int J Biochem Cell Biol.* 1998;30:293–8.
- Dewidar B, Meyer C, Dooley S, Meindl-Beinker AN. TGF-beta in hepatic stellate cell activation and liver fibrogenesis—updated 2019. *Cells.* 2019;8(11):1419.
- Dong X, Zhao B, Iacob RE, Zhu J, Koksal AC, Lu C, Engen JR, Springer TA. Force interacts with macromolecular structure in activation of TGF-beta. *Nature.* 2017;542:55–9.
- Fabregat I, Moreno-Caceres J, Sanchez A, Dooley S, Dewidar B, Giannelli G, Ten Dijke P. TGF-beta signalling and liver disease. *FEBS J.* 2016;283:2219–32.
- Gong H, Fan Z, Yi D, Chen J, Li Z, Guo R, Wang C, Fang W, Liu S. Histidine kinase NME1 and NME2 are involved in TGF-beta 1-induced HSC activation and CCl4-induced liver fibrosis. *J Mol Histol.* 2020;51:573–81.
- Hernandez-Gea V, Friedman SL. Pathogenesis of liver fibrosis. *Annu Rev Pathol.* 2011;6:425–56.
- Khalil N. TGF-beta: from latent to active. *Microbes Infect.* 1999;1:1255–63.
- Lichtman MK, Otero-Vinas M, Falanga V. Transforming growth factor beta (TGF-beta) isoforms in wound healing and fibrosis. *Wound Repair Regen.* 2016;24:215–22.
- Murphy-Ullrich JE, Poczatek M. Activation of latent TGF-beta by thrombospondin-1: mechanisms and physiology. *Cytokine Growth Factor Rev.* 2000;11:59–69.
- Parola M, Pinzani M. Liver fibrosis: pathophysiology, pathogenetic targets and clinical issues. *Mol Aspects Med.* 2019;65:37–55.
- Ramadori G, Saile B. Inflammation, damage repair, immune cells, and liver fibrosis: specific or nonspecific, this is the question. *Gastroenterology.* 2004;127:997–1000.
- Ribeiro SM, Poczatek M, Schultz-Cherry S, Villain M, Murphy-Ullrich JE. The activation sequence of thrombospondin-1 interacts with the latency-associated peptide to regulate activation of latent transforming growth factor-beta. *J Biol Chem.* 1999;274:13586–93.

- Robertson IB, Horiguchi M, Zilberberg L, Dabovic B, Hadjiolova K, Rifkin DB. Latent TGF-beta-binding proteins. *Matrix Biol.* 2015;47:44–53.
- Saharinen J, Hyytiainen M, Taipale J, Keski-Oja J. Latent transforming growth factor-beta binding proteins (LTBPs)—structural extracellular matrix proteins for targeting TGF-beta action. *Cytokine Growth Factor Rev.* 1999;10:99–117.
- Shi M, Zhu J, Wang R, Chen X, Mi L, Walz T, Springer TA. Latent TGF-beta structure and activation. *Nature.* 2011;474:343–9.
- Shrestha N, Chand L, Han MK, Lee SO, Kim CY, Jeong YJ. Glutamine inhibits CCl4 induced liver fibrosis in mice and TGF-beta1 mediated epithelial-mesenchymal transition in mouse hepatocytes. *Food Chem Toxicol.* 2016;93:129–37.
- Song X, Qiu Y, Shi J, Li L, Yuan X, Wu D, Chu Y. Prokaryotic expression, purification and evaluation of anti-cardiac fibrosis activity of recombinant TGF-beta latency associated peptide. *PeerJ.* 2022;10: e12797.
- Stachowski TR, Snell ME, Snell EH. Structural insights into conformational switching in latency-associated peptide between transforming growth factor beta-1 bound and unbound states. *IUCrJ.* 2020;7:238–52.
- Su TH, Kao JH, Liu CJ. Molecular mechanism and treatment of viral hepatitis-related liver fibrosis. *Int J Mol Sci.* 2014;15:10578–604.
- Tanwar S, Rhodes F, Srivastava A, Trembling PM, Rosenberg WM. Inflammation and fibrosis in chronic liver diseases including non-alcoholic fatty liver disease and hepatitis C. *World J Gastroenterol.* 2020;26:109–33.
- Theocharis AD, Manou D, Karamanos NK. The extracellular matrix as a multi-tasking player in disease. *FEBS J.* 2019;286:2830–69.
- Thomas GJ, Hart IR, Speight PM, Marshall JF. Binding of TGF-beta1 latency-associated peptide (LAP) to alpha(v)beta6 integrin modulates behaviour of squamous carcinoma cells. *Br J Cancer.* 2002;87:859–67.
- Unsal V, Cicek M, Sabancilar I. Toxicity of carbon tetrachloride, free radicals and role of antioxidants. *Rev Environ Health.* 2021;36:279–95.
- Xu F, Liu C, Zhou D, Zhang L. TGF-beta/SMAD pathway and its regulation in hepatic fibrosis. *J Histochem Cytochem.* 2016;64:157–67.
- Ye Q, Zhou Y, Zhao C, Xu L, Ping J. Salidroside inhibits CCl4-induced liver fibrosis in mice by reducing activation and migration of HSC induced by liver sinusoidal endothelial cell-derived exosomal SphK1. *Front Pharmacol.* 2021;12: 677810.
- Zhang CY, Yuan WG, He P, Lei JH, Wang CX. Liver fibrosis and hepatic stellate cells: etiology, pathological hallmarks and therapeutic targets. *World J Gastroenterol.* 2016;22:10512–22.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

