

REVIEW

Open Access



FADD as a key molecular player in cancer progression

Ying Liu^{1,2*} , Xiaoge Li², Xuehao Zhou², Jianxun Wang² and Xiang Ao^{1,2}

Abstract

Cancer is a leading disease-related cause of death worldwide. Despite advances in therapeutic interventions, cancer remains a major global public health problem. Cancer pathogenesis is extremely intricate and largely unknown. Fas-associated protein with death domain (FADD) was initially identified as an adaptor protein for death receptor-mediated extrinsic apoptosis. Recent evidence suggests that FADD plays a vital role in non-apoptotic cellular processes, such as proliferation, autophagy, and necroptosis. FADD expression and activity of are modulated by a complicated network of processes, such as DNA methylation, non-coding RNA, and post-translational modification. FADD dysregulation has been shown to be closely associated with the pathogenesis of numerous types of cancer. However, the detailed mechanisms of FADD dysregulation involved in cancer progression are still not fully understood. This review mainly summarizes recent findings on the structure, functions, and regulatory mechanisms of FADD and focuses on its role in cancer progression. The clinical implications of FADD as a biomarker and therapeutic target for cancer patients are also discussed. The information reviewed herein may expand researchers' understanding of FADD and contribute to the development of FADD-based therapeutic strategies for cancer patients.

Keywords FADD, Inflammation, Drug resistance, Biomarker, Therapeutic target

Background

Cancer, one of the most prevalent malignant diseases, is the second leading cause of death around the world (Liu et al. 2022f). According to the latest data from the GLOBOCAN project and the United Nations, there will be approximately 4,820,000 and 2,370,000 new cancer diagnoses and almost 3,210,000 and 640,000 cancer-related deaths in China and the United States in 2022, respectively (Xia et al. 2022). In recent years, the incidence and

mortality of cancer have gradually increased due to rapid population aging and the accumulated effects of risk factor exposure. Despite significant advances in therapeutic interventions, cancer remains a major threat to human health due to its highly lethal features and high recurrence rate (Wang et al. 2022; Zuo et al. 2022). Clinically, the majority of cancer patients are routinely diagnosed at a relatively advanced stage, with a poor five-year survival (Wang et al. 2021; Zhang et al. 2022). This is mainly caused by limited knowledge regarding the pathogenesis of cancer, lack of effective diagnosis and treatment methods, and the development of therapy resistance. Therefore, there is great urgency in clarifying the mechanisms involved in cancer progression as well as identifying effective therapeutic targets and screening diagnostic and prognostic biomarkers with high specificity and sensitivity in cancer treatment.

*Correspondence:

Ying Liu

yingliu@qdu.edu.cn; liuying_hero@163.com

¹Institute for Translational Medicine, The Affiliated Hospital of Qingdao University, Qingdao Medical College, Qingdao University, 266021 Qingdao, China

²School of Basic Medical Sciences, Qingdao Medical College, Qingdao University, 266071 Qingdao, China



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.




Fig. 1 Protein structure and PTM sites of human FADD. The structural domains are presented in the bar. The well-known PTM sites are indicated at the corresponding position. The proteins interacting with FADD are exhibited above the lines at the corresponding domains. PTM, post-translational modification; FADD, Fas-associated protein with death domain; DED, ; DD, death effector domain; NLS, nuclear localization signal; NES, nuclear export signal; cFLIP, cellular FLICE (caspase 8)-inhibitory protein; DR, death receptor; RIPK1, receptor-interacting serine/threonine protein kinase 1; ATG5, autophagy related-protein 5

Fas-associated protein with death domain (FADD), also known as MORT-1, is an evolutionarily conserved death domain (DD)-containing cytoplasmic adaptor protein and is widely expressed in various species (Mouasni and Tourneur 2018). FADD was initially reported as an adaptor molecule that bridges activated death receptors (DRs) with initiator caspase to mediate DR-induced extrinsic apoptosis (Lee et al. 2012b; Sprick et al. 2000). It is also a key modulator in non-apoptotic cellular processes, such as proliferation, cell cycle, and autophagy (Alappat et al. 2003; Antunovic et al. 2019; Imtiyaz et al. 2009). FADD has been shown to play a vital role in a variety of physiological and pathological processes, including embryonic development, innate immunity, and inflammation (Lee et al. 2012b). Therefore, the dysregulation of FADD is closely correlated with many diseases, particularly cancer (Debnath et al. 2022; Lin et al. 2020; Meer et al. 2022). The aberrant expression of FADD has been observed in multiple solid tumors, such as glioma, non-small cell lung cancer (NSCLC), and hepatocellular carcinoma (HCC) (Chen et al. 2021; Han et al. 2020; Harari-Steinfeld et al. 2021). FADD dysregulation contributes to cancer progression by affecting many aspects of cancer cell behavior, including proliferation, apoptosis, cell cycle, autophagy, inflammation, and drug resistance (Marin-Rubio et al. 2019b; Mrkvova et al. 2021). However, the exact FADD mechanisms involved in cancer progression are still not fully understood. In addition, due to its crucial roles and differential expression patterns, FADD has exhibited great potential as a valuable therapeutic target and biomarker for cancer diagnosis and treatment prognosis.

In this review, we mainly summarize recent findings on the structure, functions, and regulatory mechanisms of FADD, focus on its mechanisms of action in cancer

progression, and highlight its clinical implications in cancer treatment. We also discuss future research directions aimed at designing FADD-based therapeutic strategies for cancer patients.

Structure and functions of FADD

Structural features of FADD

The *FADD* gene is mapped to the human chromosomal band 11q13 and encodes a functional FADD protein (about 23 kDa) composed of 208 amino acids. The structure of FADD is evolutionarily conserved across species and contains two main domains, the C-terminal DD and the N-terminal death effector domain (DED) (Fig. 1). The DD (amino acids 97 to 181) of FADD mainly mediates its binding to the DRs (e.g., Fas and DR3-5) of the tumor necrosis factor (TNF) receptors superfamily to transmit apoptosis-initiating signals (Marin-Rubio et al. 2019b). Moreover, the DD mediates the interaction of FADD with receptor-interacting protein kinase (RIPK) 1 and autophagy related-protein 5 (ATG5), thereby inducing necrosis and autophagy (Pyo et al. 2005; Vanden Berghe et al. 2004). The DED also contains a nuclear localization signal (NLS) and a nuclear export signal (NES) that determine the translocation of FADD between the nucleus and cytoplasm (Gomez-Angelats and Cidlowski 2003). The DED (amino acids 3 to 81) is primarily responsible for the interaction of FADD with caspases (e.g., caspase-8 and 10) to form the death-inducing signal complex (DISC) (Bhojani et al. 2005). The DED of FADD could also mediate its binding to the cellular FLICE (caspase 8)-inhibitory protein (cFLIP) in promoting cellular proliferation (Vesely et al. 2009). Furthermore, the DED was found to mediate the interaction of FADD with p45 to diminish Fas-FADD-mediated death signaling (Sung et al. 2013).

Functions of FADD

FADD is a well-studied cytoplasmic adaptor protein that transmits DR-mediated extrinsic apoptotic signals (Zhou et al. 2022b). It can act as a bridge to interact with DRs and caspases-8/10 via its DD and DED, respectively, thereby forming an active DISC. In the DISC, caspases-8/10 are activated and then initiate the caspase cascade by a procaspase-3 or BID cleavage, eventually inducing apoptosis (Zhang et al. 2008). FADD is also involved in other forms of cell death. It has been identified as a negative regulator of necroptosis, a passive and traumatic death mediated by a RIPK1/RIPK3/mixed lineage kinase domain-like protein (MLKL) axis. FADD reduces the activity of MLKL by inducing a cleavage of the RIPK1/RIPK3 kinase domain by recruiting caspase-8/cFLIP, finally resulting in the inhibition of necroptosis (Lin et al. 1999; Osborn et al. 2010). Furthermore, FADD also plays a role in autophagic cell death and has been found to mediate the promotion of ATG5 on the autophagic death of HeLa cells triggered by interferon (IFN)- γ . FADD knockdown in HeLa cells suppresses ATG5-mediated cell death, leading to a switch from apoptotic signaling to necroptosis (Pyo et al. 2005). Some evidence suggests that FADD is involved in the regulation of several cellular processes, including cell proliferation, survival, and cell cycle (Hollomon et al. 2020; Wang et al. 2017). In addition, it has emerged as a component of various signalosomes, such as necrosome, innateosome, and inflammasome, strongly indicating the essential nature of its role in innate immunity and inflammation.

Structural biology is a hot research field in science that deeply uncovers the biomolecule structures and their structure-function relationships (Mihasan et al. 2019). With the rapid development of FADD structural biology, its functional characteristics are being elucidated from a higher-level vision in recent years. Cryogenic electron microscopy (cryo-EM) is a mainstream tool in structural biology, aimed at efficiently determining high-resolution structure of protein (Liu et al. 2022 g). Fox et al. analyzed the structure of FADD/Caspase-8 complexes using cryo-EM. They found that FADD-nucleated tandem DED (tDED) helical filament orientated the procaspase-8 catalytic domains, thereby activating them through anti-parallel dimerization. The resultant catalytically active FADD/Caspase-8 complex triggered cellular apoptosis. Moreover, c-FLIPS could be recruited into this complex and decrease Caspase-8 activity by altering tDED triple helix architecture (Fox et al. 2021). Eberstadt et al. analyzed the three-dimensional solution structure of a soluble, biologically active mutant of FADD DED using Nuclear Magnetic Resonance, a powerful method for detecting protein structure, interaction, and dynamics at atomic resolution. The structure consisted of six antiparallel, amphipathic α -helices, and the hydrophobic region

of FADD DED involving F25 was vital for its binding to FLICE and inducing apoptosis (Eberstadt et al. 1998).

Collectively, these studies strongly suggest that FADD is a key regulators in a variety of physiological and pathological processes. Currently, an increasing number of studies suggest that FADD dysregulation contributes to cancer progression. However, further research is required to elucidate the underlying mechanisms of FADD involved in cancer progression, which may be of great benefit to the development of FADD-based therapeutic strategies for cancer patients.

Molecular mechanisms of FADD regulation

FADD expression and activity are modulated by distinct mechanisms, including genetic and chromosomal alterations, DNA methylation, transcription factors (TFs), non-coding RNAs (ncRNAs), and post-translational modifications (PTMs) (Fig. 2). Herein, we present a summary of the main modes of FADD regulation, focusing specifically on the mechanisms of FADD regulation in cancer.

Genetic and chromosomal alterations contribute to FADD regulation

DNA amplification is one of the major events resulting in high FADD expression. FADD-containing DNA amplification of the 11q13 region has been observed in many types of cancer, including breast cancer (BC), bladder cancer (BCa), laryngeal/pharyngeal cancer, oral squamous cell carcinoma (OSCC), lung squamous cell carcinoma (LUSC), ovarian cancer (OC), esophageal squamous cell carcinoma (ESCC), and head and neck squamous cell carcinoma (HNSCC) (Brown et al. 2008; Chien et al. 2016; Eytan et al. 2016; Gibcus et al. 2007; Katoh and Katoh 2005; Petersen et al. 2000; Qin et al. 2016; Zhou et al. 2022a). Among them, the amplification of the *FADD* gene is most prominent in ESCC and HNSCC, with approximately 60% and 30% of patient samples harboring the amplification, respectively (Marin-Rubio et al. 2019b). Allelic losses or chromosomal deletions may result in low FADD expression. Loss of heterozygosity and deletions encompassing the *FADD* gene have been observed in 25% of cervical cancer (CC) samples (Srivatsan et al. 2002). In addition, mutations and polymorphisms in *FADD* gene have also been reported (Soung et al. 2004; Villa-Morales et al. 2010), indicating that they may also alter FADD expression and activity.

FADD regulation at the transcription level

DNA methylation is a classical mechanism that regulates gene expression. Promoter hypermethylation suppresses gene expression at the transcriptional level, whereas hypomethylation plays the opposite role (Liu et




Fig. 2 Regulation of FADD. FADD expression and activity are regulated by distinct mechanisms, including genetic and chromosomal alterations, DNA methylation, TFs, ncRNAs, and PTMs. (A–F) HIF-1 α , allelic losses or chromosomal deletions, DNA amplification, BRCA1, gene mutations or polymorphisms and DNA methylation alter FADD expression and activity at the transcription level. (G) FIST/HIPK3, PLK1, and CK1 α facilitate the translocation of FADD from the cytoplasm to the nucleus by mediating its phosphorylation at Ser194. (H) CK2 promote the nuclear localization by mediating its phosphorylation at Ser200. (I) AURA phosphorylates FADD at Ser203 in response to Taxol exposure. (J) LUBAC, MKRN1, and CHIP inhibit the formation of DISC by facilitating the degradation of FADD in an ubiquitin-proteasome dependent manner. (K) NleB1 promoted the GlcNAcylation of FADD at Arg117, thereby antagonizing the DR-induced apoptosis. (L) SUMOylation of FADD facilitates the recruitment of Drp1 onto the mitochondria by enhancing its binding to Drp1, resulting in mitochondrial-fragmentation-associated necrosis. (M) MiRNA promotes the degradation of FADD mRNA by directly targeting its 3'UTR. (N) H19 downregulates FADD by sponging miR-675. (O) Circ 001418 increased FADD levels by targeting miR-1297

al. 2019). *FADD* promoter hypermethylation has been found in some diseases, such as apical periodontitis, myelodysplastic syndrome, and OSCC. Correspondingly, low *FADD* expression has been observed in these diseases (Saberi et al. 2014; Wichniewski et al. 2019; Zhao et al. 2019). Moreover, *FADD* is a downstream target gene of several TFs, such as the hypoxia-inducible factor-1 α (HIF-1 α) and the BC susceptibility gene 1 (BRCA1). Hindryckx et al. discovered that *FADD* promoter contained a hypoxia-responsive element (HRE). Functional analysis revealed that HIF-1 α inhibited the transcriptional activity of *FADD* gene in colon cancer cells by directly binding to HREs in its promoter. Consistent with this, the deletion of the core nucleotides in the HRE of *FADD* promoter resulted in the loss of HIF-1 α -mediated transcriptional

activity suppression (Hindryckx et al. 2010). In another study, Nguyen et al. showed that BRCA1 overexpression in BRCA1-deficient BC cells significantly upregulated *FADD* expression by directly binding to *FADD* promoter. By contrast, BRCA1 knockdown efficiently decreased *FADD* expression at both the protein and messenger RNA (mRNA) levels (Nguyen et al. 2018).

Post-transcriptional regulation of FADD by ncRNAs

NcRNAs are a group of regulatory RNA molecules of different sizes, structures, and biological functions, with limited or no protein-coding capacity (Wang et al. 2020; Wen et al. 2021; Zhang et al. 2021c). They play a vital role in almost all biological processes by modulating the expression of specific gene targets (Zhang et al. 2020a, c,

Table 1 NcRNAs targeting FADD in disease

Disease types	ncRNAs	Key messages	References
Cancer	miR-128a	MiR-128a overexpression enhanced Fas-resistance on Jurkat cells by directly targeting FADD. MiR-128a knockdown sensitized Jurkat/R cells to the Fas-induced apoptosis through releasing FADD expression.	Yamada et al. (Yamada et al. 2014)
	miR-633	MiR-633 inhibited FADD expression by directly targeting its 3'UTR, leading to the enhancement of DOX/CDDP resistance in GC.	Pang et al. (Pang et al. 2019)
	H19	Overexpression of miR-675 derived from lncRNA <i>H19</i> promoted liver necroptosis and cell death in HCC cells by targeting FADD.	Harari-Steinfeld et al. (Harari-Steinfeld et al. 2021)
	miR-675	Overexpression of H19 and miR-675 facilitated proliferation and suppressed apoptosis in GC cells by inhibiting the caspase cleavage cascades (caspase 8 and caspase 3) via downregulating FADD.	Yan et al. (Yan et al. 2017)
		Overexpression of H19 increased PTK2 levels by targeting miR-138 and decreased FADD levels by targeting miR-675, leading to the inhibition of cell apoptosis and promotion of hepatoblastoma carcinogenesis.	Ge et al. (Ge et al. 2019)
	circ 001418	Overexpression of circ 001418 facilitated cell proliferation and transfer, and reduced apoptosis by upregulating EphA2 and cytochrome c protein expression, and downregulating FADD via sponging miR-1297.	Peng et al. (Peng et al. 2021)
	miR-1297		
Myocardial infarction	miR-103	Inhibition of miR-103 significantly improved the myocardial infarction of mice by releasing FADD.	Zaafan et al. (Zaafan and Abdelhamid 2021)
	H19 miR-103/107	FADD was upregulated by H19 via sponging miR-103/107. FADD upregulation promoted necrosis and improved myocardial infarction by influencing the formation of RIPK1 and RIPK3 complexes.	Wang et al. (Wang et al. 2015)
Atherosclerosis	miR-155	MiR-155 reduced the macrophage apoptosis by downregulating FADD via directly binding to its 3'-UTR region.	Zhu et al. (Zhu et al. 2013)
Intervertebral disc degeneration	miR-155	MiR-155 facilitated Fas-mediated apoptosis of nucleus pulposus cells by targeting FADD and caspase-3 via directly binding to their 3'-UTRs.	Wang et al. (Wang et al. 2011a)

2021b). NcRNAs can be divided into distinct categories, including microRNAs (miRNAs), long non-coding RNAs (lncRNAs), circular RNAs (circRNAs), and piwi-interacting RNAs (piRNAs), based broadly on their size and function (Chang et al. 2022a, b; Jia et al. 2022; Liu et al. 2021a, b; Zhang et al. 2021a). Recent studies have demonstrated that FADD expression is regulated by a series of ncRNAs in a variety of diseases (Table 1).

MiRNAs are short single-stranded ncRNAs with 18–25 nucleotides in length (Zhang et al. 2018). Their canonical role is to suppress the expression of specific target genes by directly binding to the 3'-untranslated region (3'UTR) of the mRNA (Zhang et al. 2020b; Zhou et al. 2022c). FADD is a target of miRNAs. In our previous work, we showed that high miR-633 expression was closely associated with FADD loss in gastric cancer (GC) tissue and cell lines. The knockdown of miR-633 significantly upregulated FADD in GC cells. Functional analysis revealed that miR-633 significantly decreased FADD levels by directly targeting the 3'UTR of its mRNA, resulting in the enhancement of doxorubicin/cisplatin (DOX/CDDP) resistance in GC (Pang et al. 2019). In another study, we found that miR-103/107 contributed to necrotic cell death of H9c2 cardiomyocytes by directly targeting FADD. Consistent with this, miR-103/107 knockdown blocked necrosis in a cellular model as well as myocardial

infarction in a mouse ischemia/reperfusion model (Wang et al. 2015). Furthermore, FADD was identified as a direct target of miR-155 in nucleus pulposus (NP) cells. MiR-155 inhibited FADD expression by directly targeting its 3'UTR, thereby promoting Fas-mediated apoptosis in human intervertebral disc degeneration (Wang et al. 2011a). In addition, FADD expression was modulated by a series of miRNAs, such as miR-675, miR-7a, and miR-128a, in several types of cancer (Ge et al. 2019; Liu et al. 2016; Yamada et al. 2014). lncRNAs and circRNAs are crucial regulators of FADD. For instance, lncRNA *H19* was found to decrease FADD levels by targeting miR-675 in GC, HCC, and liver necroptosis (Ge et al. 2019; Harari-Steinfeld et al. 2021; Yan et al. 2017). Circ 001418 upregulated FADD by targeting miR-1297 in BCa (Peng et al. 2021).

Taken together, these studies strongly indicate that FADD is tightly controlled by an ncRNA-based network. The dysregulation of FADD induced by this network may contribute to the development many diseases, particularly cancer. Therefore, a complete clarification of the ncRNA-based regulatory network of FADD may be highly beneficial to the development of effective therapeutic strategies for cancer patients.

Importance of PTMs in modulating FADD

PTMs are dynamic and reversible processes occurring in protein during or after their synthesis. They play a vital role in modulating protein functions such as subcellular localization, stability, and activity (Liu et al. 2022c; Liu et al. 2022 h). Studies have increasingly demonstrated that FADD can be modified by various PTMs, such as phosphorylation, ubiquitination, SUMOylation, and GlcNAcylation. FADD was found to be a substrate of multiple kinases, including FIST/HIPK3 kinase, casein kinase (CK) I α , polo kinase (PLK)1, Aurora-A kinase (AURA), and CK2 (Alappat et al. 2005; Jang et al. 2011a, b; Rochat-Steiner et al. 2000; Sheikh and Huang 2003; Vilmont et al. 2015). Distinct kinases can phosphorylate FADD at diverse sites and generate a range of cellular effects, and FADD can be phosphorylated by FIST/HIPK3, CKI α , and PLK1 at Ser194 (Alappat et al. 2005; Jang et al. 2011b; Rochat-Steiner et al. 2000). The phosphorylation of this site mediated the translocation of FADD between the cytoplasm and the nucleus (Sheikh and Huang 2003). AURA directly phosphorylated FADD at Ser203 in response to Taxol treatment (Jang et al. 2011a). Moreover, FADD was phosphorylated by CK2 at Ser200, which was required for its nuclear localization (Vilmont et al. 2015). Ubiquitination is a crucial PTM that mediates the degradation of proteins in a proteasome-dependent manner. Currently, several E3 ubiquitin ligases for FADD have been identified, including the makorin ring finger protein 1 (MKRN1), the C terminus HSC70-interacting protein (CHIP), and the linear ubiquitin chain assembly complex (LUBAC) (Goto and Tokunaga 2017; Lee et al. 2012a; Seo et al. 2018). Lee et al. discovered that MKRN1 promoted the ubiquitin-dependent degradation of FADD. MKRN1 knockdown enhanced FADD stability and facilitated DISC formation, thereby causing hypersensitivity to extrinsic apoptosis (Lee et al. 2012a). Seo et al. revealed that CHIP inhibited DISC formation and apoptosis in HeLa cells by inducing the polyubiquitylation of FADD at Lys149 and 153. Consistent with this, CHIP deficient in E3 ligase activity failed to suppress the function and ubiquitylation of FADD (Seo et al. 2018). Moreover, FADD was a substrate of LUBAC, which mediated its linear ubiquitination upon TNF α stimulation in T-cell leukemia Jurkat cells (Goto and Tokunaga 2017).

SUMOylation is a type of ubiquitination-like PTM that plays a vital role in the modulation of protein function (Fan et al. 2022). FADD was found to be modified by SUMO2 at Lys120/125/149 during intrinsic necrosis of HeLa cells. SUMOylation promoted dynamin-related protein 1 (Drp1) recruitment onto the mitochondria by enhancing the binding of FADD to Drp1, resulting in Drp1-mediated mitochondrial fragmentation. Consistent with this, SUMO-defective FADD expression in murine embryonic fibroblasts (MEFs) blocked

mitochondrial-fragmentation-associated necrosis (Choi et al. 2017). In addition, FADD is a substrate of NleB1, an arginine glycosyltransferase that modifies conserved arginine residues in DD-containing host proteins with N-acetylglucosamine (GlcNAc). NleB1 promoted the GlcNAcylation of FADD at Arg117 in HeLa cells during bacterial infection, thereby antagonizing the DR-induced apoptosis of infected cells (Scott et al. 2017). Collectively, these findings suggest that FADD activity is regulated by various PTMs. Thus in-depth investigations are required to further explore the PTM network involved in FADD regulation.

Implications of FADD in cancer progression

FADD was deficient in certain types of cancer, such as thymic lymphoma, acute myeloid leukemia (AML), and glioblastoma (GBM) (Newton et al. 2000; Tourneur et al. 2004; Wang et al. 2017). However, high FADD expression was also observed in multiple cancer types, including pancreatic cancer (PC), laryngeal/pharyngeal cancer, OSCC, HNSCC, and BC (Chien et al. 2016; Gibcus et al. 2007; Gonzalez-Moles et al. 2020; Zhang et al. 2017; Zhou et al. 2022a). These findings strongly suggest that FADD may play a dual role in cancer progression by acting as a tumor suppressor or oncoprotein. However, the exact mechanisms are yet to be fully understood. In the following section, we summarize the recent advances regarding the role of FADD in cancer progression (Table 2).

FADD and cancer-related signaling pathways

The crosstalk of FADD with cancer-related signaling pathways, such as mitogen-activated protein kinase (MAPK) and nuclear factor- κ B (NF- κ B) signaling pathways, is considered a main mechanism through which FADD participates in the modulation of cancer progression. The MAPK signaling pathway is a conserved oncogenic pathway, and its aberrant activation has been shown to facilitate cancer progression by regulating multiple cellular processes, such as proliferation, apoptosis, and drug resistance (Liu et al. 2022 h). MAPKs mainly consist of three members of the c-Jun NH2 terminal kinase (JNK), extracellular stress regulated kinase (ERK), and p38 (Kciuk et al. 2022). Shimada et al. explored the effect of FADD phosphorylation mediated by JNK1 on drug sensitivity in prostate cancer (PCa) using cell models. Western blotting (WB) and flow cytometry (FCM) assays were performed. The results showed that FADD was phosphorylated by JNK1 in LNCaP and DU145 cells treated with paclitaxel (PTX), resulting in the enhancement of PTX sensitivity in PCa. Mechanistically, PTX administration increased MEK kinase 1 (MEKK1) levels in PCa cells, leading to the activation of JNK. Next, FADD was phosphorylated by JNK1 at Ser194, which further upregulated MEKK1 and enhanced downstream

Table 2 Functional roles of FADD in different types of cancer

Cancer types	Key messages	References
BC	High expression or amplification of <i>FADD</i> was negatively associated with the abundance of CD4+ T cells and dendritic cell infiltration in BC ($p < 0.05$). High expression of <i>FADD</i> mRNA significantly associated with RFS BC in patients ($p < 0.05$). High expression of <i>FADD</i> was frequently detected in luminal B and high-grade BC with shorter metastasis-free survival times ($p < 0.05$).	Zhou et al. (Zhou et al. 2022a)
	Increase copy number and mRNA overexpression of <i>FADD</i> are frequently observed in primary BC.	Callegari et al. (Callegari et al. 2016)
	<i>FADD</i> knockdown promoted autophagy of BC cells by downregulating Rheb and inhibiting mTORC1 activity.	He et al. (He et al. 2016)
	<i>FADD</i> expression was significantly associated with T stage ($P = 0.046$). The combination of <i>FADD</i> , <i>PPF1A1</i> , and <i>TMEM16A</i> genes was significantly correlated with perineural invasion ($p = 0.022$) and DFS ($p = 0.034$).	Choi et al. (Choi et al. 2014)
	<i>MKRN1</i> mediated the ubiquitination and proteasomal degradation of <i>FADD</i> . The stabilization of <i>FADD</i> facilitated extrinsic apoptosis by promoting caspase-8 and caspase-3 cleavage in BC cells. <i>FADD</i> participated in the modulation of necroptosis upon caspase inhibition.	Lee et al. (Lee et al. 2012a)
JNK-mediated phosphorylation of <i>FADD</i> inhibited cell growth and metastasis through G2/M arrest.	Matsuyoshi et al. (Matsuyoshi et al. 2006)	
GC	<i>FADD</i> was negatively regulated by miR-633, and its upregulation enhanced DOX/DDP induced apoptosis in GC cells.	Pang et al. (Pang et al. 2019)
	<i>FADD</i> expression was negatively associated with H19 and miR-675 expression in GC. <i>FADD</i> mediated the promotion of H19/miR-675 axis on GC progression.	Yan et al. (Yan et al. 2017)
	The expression of p- <i>FADD</i> in GC was heterogenous in its location, whereas its expression in normal gastric cells was uniform nuclear expression.	Yoo et al. (Yoo et al. 2007)
NSCLC	NSCLC patients with overexpression of <i>FADD</i> had lower OS rates ($p = 0.033$). The upregulation of <i>FADD</i> was an independent poor prognostic biomarker for patients with surgically resected NSCLC ($p = 0.027$).	Chen et al. (Chen et al. 2021)
	High <i>FADD</i> expression predicted poor prognosis of NSCLC patients. The ubiquitin ligase SPOP promoted <i>FADD</i> degradation by directly binding to it. <i>FADD</i> mediated the inhibition of SPOP on NF- κ B activity and its target genes expression.	Luo et al. (Luo et al. 2018)
	<i>FADD</i> was specifically downregulated in NSCLC cells, and <i>FADD</i> loss was significantly associated with the presence of extracellular <i>FADD</i> . NSCLC tissues released significantly more <i>FADD</i> than non-tumoral tissue ($P = 0.000003$). The release of <i>FADD</i> increased significantly with the cancer stage, and was associated with both early and late steps of the metastasis process.	Cimino et al. (Cimino et al. 2012)
PCa	PCa patients with a greater positive p- <i>FADD</i> rate had a significantly lower biochemical recurrence rate than those with a lower positive p- <i>FADD</i> rate ($p < 0.001$). The positive p- <i>FADD</i> rate was negatively associated with Gleason score.	Ikeda et al. (Ikeda et al. 2013)
	The overexpression of dephosphorylated <i>FADD</i> (S194A) promoted proliferation and invasion of PCa cells by enhancing hTERT expression and telomerase activity. PCa patients expressing low levels of p- <i>FADD</i> had significantly higher rates of biochemical recurrence than those with high p- <i>FADD</i> expression ($p < 0.001$).	Matsumura et al. (Matsumura et al. 2009)
	The overexpression of phosphorylated <i>FADD</i> resulted in G2/M cell-cycle arrest, whereas the non-phosphorylated <i>FADD</i> overexpression resulted in cell cycle progression and enhanced colony-forming activity in PCa cells.	Shimada et al. (Shimada et al. 2005)
	<i>FADD</i> was phosphorylated by JNK at Ser194 in PCa cells treated with PTX. Phosphorylated <i>FADD</i> enhanced the synergistic effects of paclitaxel on anticancer drug-induced apoptosis in PCa cells.	Shimada et al. (Shimada et al. 2004)
	<i>FADD</i> mediated the effect of PTEN in promoting drug-induced apoptosis by facilitating caspase-8 activation and BID cleavage in PCa.	Yuan et al. (Yuan and Whang 2002)
Leukemia	<i>FADD</i> knockdown in Jurkat cells significantly inhibited cell proliferation, enhanced sensitivity to Etoposide-induced intrinsic apoptosis and resistance to TRAIL-induced extrinsic apoptosis by triggering a metabolic shift from glycolysis to mitochondrial respiration.	Zhou et al. (Zhou et al. 2022b)
	<i>FADD</i> knockdown in Jurkat cells enhanced drug resistance, which could be partially overcome by induction of RIP1-dependent necroptosis through TNFR1 activation using combined treatment with TNF α and LCL161.	Mrkvoa et al. (Mrkvoa et al. 2021)
	S194-P- <i>FADD</i> was more stable and preferentially localized to the cell nucleus, thereby promoting proliferation in T-cell lymphoblastic lymphoma cells. <i>FADD</i> phosphorylation was a prognostic biomarker in T-cell lymphoblastic lymphoma.	Marin-Rubio et al. (Marin-Rubio et al. 2019a)
	The availability of <i>FADD</i> in the cytoplasm reduced in T-cell lymphoblastic lymphoma cells. Reduction of <i>FADD</i> phosphorylation that inversely correlates with the proliferation capacity and tumor aggressiveness. The reduction of <i>FADD</i> phosphorylation was negatively associated with the proliferation capacity and tumor aggressiveness.	Marin-Rubio et al. (Marin-Rubio et al. 2016)
	<i>FADD</i> was a direct target of miR-128a. <i>FADD</i> downregulation by miR-128a conferred Fas-resistance on Jurkat cells. <i>FADD</i> upregulation induced by miR-128a knockdown sensitized Jurkat/R cells to the Fas-mediated apoptosis.	Yamada et al. (Yamada et al. 2014)

Table 2 (continued)

Cancer types	Key messages	References
Glioma	FADD was downregulated by ATRX via the H3K27me3 enrichment, resulting in the enhancement of PARP1 stabilization in TMZ resistant glioma cells.	Han et al. (Han et al. 2020)
	FADD overexpression induced apoptosis in about 85% of malignant glioma cells. The retroviral transfer of FADD gene significantly suppressed survival in malignant glioma cells by induction of apoptosis.	Kondo et al. (Kondo et al. 1998)
GBM	FADD overexpression inhibited proliferation and induced apoptosis in human GBM cells.	Wang et al. (Wang et al. 2017)
HCC	FADD mediated the role of OTULIN in preventing the development of chronic liver inflammation and HCC.	Verboom et al. (Verboom et al. 2020)
	FADD mediated the apoptosis of hepatocytes, thereby promoting the development of hepatitis and HCC in NEMO(LPC-KO) mice	Ehlken et al. (Ehlken et al. 2014)
	FADD mediated the role of truncated RIP3 (aa 224–518) in inducing apoptosis in human hepatocellular carcinoma cells QGY-7703.	Feng et al. (Feng et al. 2006)
	FADD expression was significantly correlated with cell apoptosis in HCC ($p < 0.05$). The positive rate of FADD expression in HCC is lower than that in adjacent normal tissues ($p < 0.05$).	Sun et al. (Sun et al. 2000)
Melanoma	The degradation of FADD was inhibited by ADT-OH via downregulating MKRN1. The tumor-specific delivery of FADD combined with low-dose ADT-OH administration significantly suppressed tumor growth and induced cell apoptosis in melanoma.	Cai et al. (Cai et al. 2020)
	The overexpression of FADD or truncated FADD (1-181 aa) induced apoptosis in B16F10 cells, The truncated FADD exhibited a more potent apoptotic effect than FADD. The tumor-targeted delivery of FADD or truncated FADD inhibited tumor growth by inducing apoptosis of tumor cells via activating caspase-dependent apoptotic pathway.	Yang et al. (Yang et al. 2016)
	FADD decreased FAK expression by upregulating miR-7a, thereby promoting cell migration in melanoma.	Liu et al. (Liu et al. 2016)
OC	FADD mediated the induction of apoptosis by certain chemotherapeutic drugs in OC cells.	Milner et al. (Milner et al. 2002)
CRC	Phosphorylated FADD interacted with MT2A and co-localized with MT2A mostly to nuclei and slightly to cytoplasm. The co-expression of Phosphorylated FADD and MT2A inhibited the apoptosis and promote proliferation in CRC cells.	Marikar et al. (Marikar et al. 2016)
	The overexpression of FADD mediated by adenovirus inhibited cell growth and enhanced apoptosis of SW480 cells and suppress growth of xenografts in mice.	He et al. (He et al. 2018)
	FADD mediated apoptosis of CRC cells synergistically induced by Smac mimetic and DOX.	Yang et al. (Yang et al. 2020b)
	Apigenin upregulated FADD and induced its phosphorylation, which may resulted in cell apoptosis and inhibition of tumor growth in CRC.	Wang et al. (Wang et al. 2011b)
	FADD overexpression enhanced 5-FU sensitivity in CRC mice model. Stable overexpression of FADD significantly improved apoptosis-inducing effects of 5-FU on CRC cells.	Yin et al. (Yin et al. 2010)
HNSCC	The immunohistochemical overexpression of FADD was significantly associated with worse OS ($p < 0.001$), DSS ($p < 0.001$), DFS ($p < 0.001$), higher clinical stage ($p = 0.005$), and a large magnitude of effect with N + status ($p < 0.001$).	Gonzalez-Moles et al. (Gonzalez-Moles et al. 2020)
	High FADD expression was significantly associated with an increased rate of lymph node metastasis ($p = 0.001$), a shorter distant metastasis-free interval ($p = 0.046$).	Pattje et al. (Pattje et al. 2013)
	HNSCC patients with tumors that were strongly positive for cyclin D1 and FADD had reduced OS ($p = 0.003$ and $p < 0.001$), DSS ($p = 0.039$ and $p < 0.001$), and DFS ($p = 0.026$ and $p < 0.001$), respectively. FADD was a significant independent predictor of DSS and DFS.	Rasamny et al. (Rasamny et al. 2012)
OSCC	FADD gene copy number and protein expression are potential prognostic biomarkers and are closely associated with lymph node metastasis in OSCC patients ($p < 0.001$).	Chien et al. ((Chien et al. 2016)
Clear cell renal cell carcinoma	FADD was downregulated in clear cell renal cell carcinoma compared with that in normal tissues. FADD-mediated apoptosis may suppress carcinogenesis in clear cell renal cell carcinoma.	Xu et al. (Xu et al. 2009)
PC	FADD is required for PC cell proliferation and that it is overexpressed to varying degrees in a variety of PC cell types. FADD protected PC cells from drug-induced apoptosis, whereas FADD knockdown enhanced the sensitivity of drug-resistant cells to ADR-mediated apoptosis.	Zhang et al. (Zhang et al. 2017)

JNK1 activation, finally resulting in G2/M cell cycle arrest and apoptosis of PCa cells (Shimada et al. 2004). Tewari et al. examined the role of FADD in mediating miltefosine-induced apoptosis of glioma cells

through WB and 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2 H-tetrazolium, inner salt assays. They discovered that the levels of phosphorylated ERK was significantly upregulated in

U87MG and T98G cells treated with miltefosine, and activated ERK increased FADD expression, resulting in enhancement of glioma cell apoptosis induced by miltefosine. Consistent with this, the increase in FADD expression and cell apoptosis observed in U87MG and T98G cells treated with miltefosine was blocked in the presence of U0126 (an ERK inhibitor) (Tewari et al. 2008).

The NF- κ B signaling pathway plays a vital role in regulating various biological processes, including cell differentiation, proliferation, and survival, and, most importantly, immune responses and inflammation (Chaithongyot et al. 2021). The dysregulation of NF- κ B is closely associated with a variety of human diseases, particularly cancer (Liu et al. 2022a). Chaudhary et al. examined the effect of FADD on the NF- κ B signaling pathway using a BC cell model. Luciferase reporter assay showed that low FADD concentration induced the activation of the NF- κ B signaling pathway in MCF-7 cells in a time- and dose-dependent manner, with caspase 8 or its homologs mediating this process (Chaudhary et al. 2000). FADD also exhibits the effect to inhibit NF- κ B activation. Ranjan et al. performed luciferase reporter and p65 translocation assays and discovered that FADD overexpression blocked NF- κ B activation induced by TNF α and translocation of p65 from the cytoplasm to the nucleus in MCF-7 cells and colorectal cancer (CRC) cell line HCT 116 (Ranjan and Pathak 2016b). The same team chemically conjugated purified FADD protein with cell permeable TAT (transactivator of transcription) peptide in another study, to deliver in cancer cells. Luciferase reporter and quantitative real-time polymerase chain reaction (qRT-PCR) assays showed that TAT-FADD significantly reduced the activation of NF- κ B in HCT-116 cells, leading to the inhibition of anti-apoptotic genes, including *Bcl2*, *cIAPs*, *RIP1*, and *cFLIPL* (Ranjan et al. 2020). These studies suggest that FADD possesses double-edged role in regulating the NF- κ B signaling pathway. The sufficient availability and concentration of FADD may be the cause of its differential roles in the activation of the NF- κ B signaling pathway.

Taken together, these findings demonstrate that JNK1 induces G2/M cell cycle arrest and apoptosis in PCa cells by activating FADD, whereas ERK facilitates the apoptosis of glioma cells by upregulating FADD levels. FADD also exhibits double-edged effect in targeting the NF- κ B signaling pathway in different cancer types. However, these studies have limitations. For instance, only cell models are used in these studies. Animal models and clinical samples should be utilized to validate these findings. In addition, there are various other cancer-related signaling pathways, such as the PI3K/AKT, Wnt/ β -catenin, and STAT3 signaling pathways. Whether FADD can participate in cancer progression through synergistic action with these signaling pathways has not been

reported. This issue should be included in future studies. Nevertheless, fully understanding the underlying mechanisms of FADD in the modulation of cancer-related signaling pathways may provide new insights into the development of effective therapeutic strategies for cancer patients.

FADD and proliferation

One of the most remarkable hallmarks of cancer is the uncontrolled proliferation of tumor cells. The mechanisms involved in proliferation are extremely complicated and are yet to be fully understood (Liu et al. 2022e). Accumulating evidence suggests that FADD is a key regulator of cellular proliferation during cancer progression (Marin-Rubio et al. 2019a; Wang et al. 2017; Zhang et al. 2017). For example, Wang et al. performed qRT-PCR and WB assays and found that FADD was downregulated at both mRNA and protein levels in GBM tissues compared with that in normal brain tissues. Moreover, statistical analysis (SPSS 21.0 software) showed that FADD expression in stages III and IV patients was lower than those in stages I and II. Cell count kit-8 (CCK-8) analysis further revealed that FADD overexpression significantly inhibited the proliferation of the GBM cell line SHG44 (Wang et al. 2017). Zhang et al. conducted reverse transcription-quantitative polymerase chain reaction (RT-qPCR) and WB assays to examine the expression of FADD. FADD was significantly upregulated in the PC cell line Colo357 at both mRNA and protein levels. 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium (MTT) assay showed that the knockdown of FADD significantly suppressed the proliferation of Colo357 cells (Zhang et al. 2017). Moreover, the data from fluorescence resonance energy transfer, MTT, Ki67 immunostaining, immunohistochemistry (IHC) assays demonstrated that Ser194-phosphorylated FADD was localized in the nucleus of tumor cells and induced proliferation in multiple types of cancer, including T-cell lymphoblastic lymphoma, CRC, and lung cancer (Bhojani et al. 2005; Marikar et al. 2016; Marin-Rubio et al. 2019a).

FADD could also mediate the effect of ncRNAs on proliferation in cancer. Yan et al. examined the effects H19 and miR-675 on GC cell proliferation using CCK-8 and colony formation assays and discovered that the overexpression of H19 and miR-675 facilitated the proliferation of SGC-7901 cells, whereas their knockdown had the opposite effects. The authors further explored the role of H19/miR-675 axis in GC using a nude mice model. A dramatic decrease was observed in tumor volume and weight in nude mice in which LV-shH19 infected SGC-7901 cells were injected. Co-infection with LV-miR675 partially reversed this tendency. Mechanistically, H19/miR-675 suppressed the caspase cleavage cascades (caspase 8 and caspase 3) by downregulating FADD, leading

to the inhibition of the proliferation of SGC-7901 cells (Yan et al. 2017). In another study, Peng et al. explored the mechanism of circ 001418 involved in BCa cell proliferation using MTT and WB assays. Circ 001418 overexpression was found to facilitate proliferation of BCa cells by downregulating FADD and upregulating EphA2 and cytochrome c protein via sponging miR-1297 (Peng et al. 2021).

Collectively, these findings show that FADD can act as a tumor suppressor to inhibit cellular proliferation in several cancer types, such as GBM, CRC, and lung cancer. It can also facilitate proliferation in GC and BCa by serving as an oncoprotein. However, limitation still exists in these studies. For instance, almost all in vitro culture models are established using tumor cell lines. Primary cells originated from cancer patients should be used in future studies. In addition, the exact mechanisms of FADD involved in proliferation regulation are not fully elucidated. The upstream regulators and downstream effectors of FADD remain unclear. Therefore, uncovering regulatory network based on FADD involved in proliferation regulation should be an important direction in future studies.

FADD and cell cycle

The cell cycle is a ubiquitous and complex process by which a cell duplicates its contents and divides into two genetically identical daughter cells. This process is tightly controlled by various cell cycle regulatory proteins, such as cyclin proteins, cyclin dependent kinases (CDKs), and mitotic checkpoint proteins (Roy et al. 2017). Dysregulation of the cell cycle has been shown to be a fundamental mechanism underlying carcinogenesis, making cell cycle regulators promising anti-tumoral therapeutic targets (Liu et al. 2022b). An increasing amount of evidence suggests that FADD is a key cell cycle regulator during cancer progression (Bowman et al. 2015; Chen et al. 2005; Matsuyoshi et al. 2006; Shimada et al. 2005). *Cyclin D1* and *Cyclin B1* are downstream target genes of NF- κ B and are involved in regulating cell cycle process (Zhang et al. 2015). Chen et al. performed statistical analysis to evaluate the correlation between FADD and cyclin D1/B1 protein expression. They discovered that high FADD expression in lung cancer tissues was significantly associated with the overexpression of the cyclins D1 and B1. NF- κ B TransAM kit and fluorescence-activated cell sorting assays were used to examine the role of FADD and p-FADD in cell-cycle progression. The results showed that phosphorylated FADD overexpression in Jurkat cells increased NF- κ B activity and facilitated the accumulation of cells in the G2/M phase of the cell cycle (Chen et al. 2005). In another study, Bowman et al. analyzed the role of FADD phosphorylation in the regulation of cell cycle process in lung cancer. Co-immunoprecipitation (Co-IP) and mass spectrometry assays showed that CK1 α

phosphorylated FADD by directly interacting with it. Histopathology analysis of the H&E stain indicated that phosphorylated FADD mainly localized in the nucleus of lung cancer cells from a mice model. The data from WB and mass spectrometry assays further revealed that phosphorylated FADD was most abundant during the G2/M phase of the cell cycle, and it directly interacted with kinases (PLK1, AURA, and BUB1) that mediate the G2/M transition (Bowman et al. 2015). Moreover, Shimada et al. performed FCM analysis to detect the effect of phosphorylated FADD on cell cycle in primary cultures of normal epithelial cells and DU145 cells. They demonstrated that the induction of phosphorylated FADD in prostate epithelial cells resulted in cell cycle arrest at G2/M to protect against uncontrolled proliferation. FADD was more dephosphorylated in PCa cells than in normal epithelial cells, and dephosphorylated FADD promoted the cell cycle progression of PCa cells (Shimada et al. 2005). In addition, Matsuyoshi et al. conducted WB and FCM assays and discovered that FADD was phosphorylated by JNK in the ER-negative BC cell line MDA-MB-231 treated with PTX or tamoxifen. Phosphorylated FADD could induce cell cycle arrest at G2/M and inhibit cell proliferation (Matsuyoshi et al. 2006).

These findings demonstrate that FADD can be phosphorylated by a series of kinases (e.g., CKI α , PLK1, and BUB1). Phosphorylated FADD alters the expression of cell cycle regulatory proteins (e.g., Cyclin D1 and Cyclin B1) by activating NF- κ B signaling pathway, thereby mediating the G2/M transition of cancer cells. Therefore, the phosphorylation status of FADD is crucial for the transition of cancer cells at the G2/M boundary. However, research on the role of FADD in regulating the cell cycle remains limited. For example, whether FADD can modulate cell cycle progression through other signaling pathways. In addition to phosphorylation, FADD can be modified by ubiquitination, SUMOylation, and GlcNAcylation. Whether these PTMs also play a role in cell cycle regulation during cancer progression. Future studies should focus on the clarification of these issues. Collectively, fully understanding the mechanisms of FADD in cell cycle regulation would help in the precise use of FADD-based therapeutics in particular types of cancer.

FADD and apoptosis

Apoptosis induction is a major mechanism of many anti-cancer drugs in cancer treatment as apoptosis evasion results in therapy resistance for cancer patients (Singh and Lim 2022). Complex signaling pathways and a variety of regulators are involved in cell apoptosis during cancer progression. Thus, fully elucidating the underlying mechanisms involved in apoptosis may provide novel insights on the treatment of cancer. FADD is a well-known adaptor protein for DR-mediated extrinsic

apoptosis (Zhou et al. 2022b). A large number of studies have shown that FADD is involved in cancer progression by regulating apoptosis (Hollomon et al. 2020; Mrkvořa et al. 2021; Wang et al. 2017). For instance, Hollomorn et al. performed trypan blue exclusion and WB assays to examine the effect of FADD on TNF α -induced cell death. They discovered that FADD knockdown facilitated cell apoptosis in osteosarcoma LM7 or SaOS2 cells treated with TNF α . Further analysis revealed that the inhibition of NF- κ B in wildtype LM7 cells significantly increased TNF α -induced apoptosis to similar levels observed in the FADD knockdown of LM7 cells, indicating that FADD plays a role in NF- κ B pro-survival signaling (Hollomon et al. 2020). Wang et al. conducted qRT-PCR and WB to detect FADD expression and found that FADD was downregulated in GBM tissues and cell lines SC189, u251, and SHG44. The data from FCM analysis demonstrated that FADD overexpression significantly increased the apoptosis rate of SHG44 cells. Moreover, FADD-overexpressed GBM cells showed increased expressions of FADD, caspase-8, and Bax and decreased expressions of Bcl-2 (Wang et al. 2017). Mrkvořa et al. identified molecules involved in cell death signaling pathways triggered by conventional anti-cancer drugs using Jurkat cell model with for necroptosis and/or apoptosis genes, created by the CRISPR-Cas9 editing system. The data from FCM analysis demonstrated that FADD was indispensable to the apoptosis induced by anti-cancer drugs (e.g., DOX and etoposide) in Jurkat cells. Consistent with this, Jurkat cells with knocked out FADD showed resistance to these drugs (Mrkvořa et al. 2021). Furthermore, He et al. constructed a recombinant adenovirus containing FADD gene and explored the effect of FADD on cell growth and apoptosis using colon cancer cell and mice models. Results showed that FADD overexpression suppressed cell growth and enhance apoptosis in SW480 cells in vitro. Consistence with this, its overexpression also inhibited the growth of subcutaneous xenografts in the nude mice (He et al. 2018).

Taken together, these studies demonstrate that FADD is a crucial regulator of apoptosis during cancer progression. Apoptosis can be classified into two categories: the extrinsic pathways mediated by DRs and intrinsic pathways mediated by mitochondria. FADD has been shown to be a key component for DR-mediated extrinsic apoptosis. In several studies, FADD was found to impair mitochondrial integrity and facilitate mitochondrial fragmentation (Choi et al. 2017; Ranjan and Pathak 2016a), indicating that FADD may participate in regulating mitochondria-mediated intrinsic apoptosis. Therefore, whether FADD is involved in the regulation of intrinsic apoptosis during cancer progression and the underlying mechanisms involved in this process should be further investigated in future studies. Although the detailed

mechanisms of FADD involved in apoptosis are still not fully elucidated, it has exhibited great potential as a valuable target for cancer treatment.

FADD and inflammation

Chronic inflammation is a well-known characteristic of cancer that substantially contributes to cancer initiation and progression (Murata 2018). Due to the high genetic stability of the cells responsible for cancer-associated inflammation, they are not subjected to the rapid emergence of drug resistance (Singh et al. 2019). Therefore, targeting inflammation is a promising strategy for cancer prevention and treatment. Accumulating evidence has pointed to the involvement of FADD in the regulation of inflammation during cancer progression. FADD participates and plays a vital role in most signalosome complexes, such as inflammasome and FADDosome (Henry and Martin 2017; Tummers et al. 2020). Ranjan et al. performed WB, RT-qPCR, and ELISA assays using colon cancer cell model to determine the role of FADD in inflammation regulation. They discovered that the delivery of FADD mediated by TAT in colon cancer HCT116 cells significantly suppressed the canonical NLRP3 inflammasome priming by downregulating NLRP3, thereby restricting the processing and secretion of pro-inflammatory IL-1 β . Moreover, Luciferase reporter and qRT-PCR assays demonstrated that TAT-FADD inhibited the activation of NF- κ B-induced proinflammatory priming in HCT116 cells (Ranjan et al. 2020). Henry et al. conducted co-IP, WB, ELISA assays using TRAIL-stimulated HeLa cell model and showed that FADD formed the FADDosome complex by directly interacting with caspase-8 and RIPK1, thereby facilitating NF- κ B activation and pro-inflammatory cytokine production (e.g., IL-6, IL-8, and MCP-1) (Henry and Martin 2017). In another study, Hartwig et al. validated these results in human lung adenocarcinoma (ADC) A549 cells. They examined secretion of proinflammatory cytokines by ELISA assay and found that that FADD mediated the promotion of TRAIL upon secretion of proinflammatory cytokines IL-8 and MCP-1 in human lung adenocarcinoma (ADC) A549 cells. Consistent with this, FADD knockdown inhibited NF- κ B activation and inflammatory cytokine production in A549 cells in response to TRAIL stimulation (Hartwig et al. 2017).

Collectively, these findings show that FADD exerts proinflammatory effects in multiple types of cancer, including CRC, CC, and ADC. However, FADD can also contribute to anti-inflammatory responses of some non-tumor cell lines (Hu et al. 2000; Schaub et al. 2000). For instance, FADD overexpression in MEFs was found to suppress LPS-induced NF- κ B activation but promote LPS-induced IFN β production (Brennan et al. 2019), which means that FADD may possess great potential

to play an anti-inflammatory role in some other cancer types. This hypothesis should be further tested in future studies. In addition, there are limitations in these studies as only cell models were used in exploring the role of FADD in inflammation. Animal models and patient samples should be applied in future studies to validate these findings. Nevertheless, in-depth investigations are required to elucidate the detailed mechanisms of FADD involved in inflammation regulation during cancer progression, which may provide new insights into the development of FADD-based therapeutic strategies for cancer patients.

FADD and drug resistance

The development of drug resistance to chemotherapy continues to be one of the important reasons for treatment failure and patient death in cancer (Liu et al. 2022d). The mechanisms underlying drug resistance mainly include tumor heterogeneity, some cellular-level changes, and genetic factors, though not fully understood (Nussinov et al. 2021). It has been reported that FADD is a crucial regulator of drug resistance during cancer progression (Han et al. 2020; Mrkvova et al. 2021; Pang et al. 2019). For instance, in our previous study, FADD was found to mediate the inhibitory effect of miR-633 on drug resistance in GC cell models. The data from WB, qRT-PCR and TUNEL assays demonstrated that FADD was downregulated by miR-633, which significantly enhanced the apoptosis of SGC-7901 cells induced by DOX/CDDP (Pang et al. 2019). Han et al. showed that FADD plays a vital role in temozolomide (TMZ)-mediated drug resistance in glioma. They performed immunofluorescence assay using glioma cell model and discovered that FADD overexpression increased temozolomide (TMZ)-induced DNA damage in glioma cells, whereas FADD knock-down decreased TMZ-induced DNA damage in ATRX knockout glioma cells. Furthermore, the data from WB, qRT-PCR, and immunofluorescence assays revealed that FADD expression was inhibited in TMZ-resistant glioma cells by ATRX by promoting the trimethylation of histone H3K27 in the *FADD* promoter region. The downregulation of FADD mediated by ATRX promoted PARP1 stabilization, resulting in the enhancement of DNA damage repair (Han et al. 2020). Moreover, Zhang et al. conducted RT-qPCR, WB, and MTT assays to analyze the effect of FADD on drug resistance in PCa cell models. They found that FADD was upregulated to varying degrees in various types of PCa cells. This results in differing levels of drug resistance in PCa cells. Furthermore, drug resistance is positively associated with FADD expression. The data from Annexin V immunostaining analysis revealed that FADD could protect PCa cells from Adriamycin-induced apoptosis (Zhang et al. 2017).

These studies strongly suggest that FADD plays a double-edged role in the development of cancer drug resistance. FADD can inhibit drug resistance in glioma cells through enhancing DNA damage induced by TMZ. It can also enhance drug resistance in GC and AML cells through suppressing apoptosis induced by chemotherapeutic agents (e.g., DOX and Adriamycin). However, the exact effects of FADD on drug resistance in cancer are still unclear. For instance, whether FADD possesses the same effect in mediating one anti-cancer drug in different cancer types. Whether FADD have the same role in mediating multiple anti-cancer drugs in one cancer. Moreover, multiple mechanisms, such as excessive drug efflux, alteration of CSC characteristics, and autophagy dysregulation, can result in the development of drug resistance in cancer. Whether FADD is involved in cancer drug resistance by regulating these processes. All these concerns should be addressed in future studies. Taken together, further studies on the mechanisms of FADD involved in drug resistance in distinct types of cancer will be of great benefit to the development of FADD-based therapeutic strategies for patients demonstrating a poor response to chemotherapy.

In summary, FADD may play the dual role of tumor suppressor or oncoprotein in cancer progression. FADD dysregulation contributes to cancer progression by influencing many aspects of cancer cell behavior, including proliferation, apoptosis, cell cycle, inflammation, and drug resistance (Fig. 3). The double-edged effect of FADD on the development of cancer drug resistance may depend on its diverse functions.

Clinical applications of FADD in cancer treatment

FADD as a prognostic biomarker in cancer

Prognostic assessment is a core link in the management of cancer patients. Accurately assessing a prognosis can help judge the efficiency of a current therapeutic strategy and adjust a proper treatment plan for cancer patients in a timely manner (Simmons et al. 2017). However, patient outcomes continue to be poor due to the lack of effective evaluation methods. Biomarkers are considered a promising tool in the early diagnosis, prognostic assessment, and treatment of cancer patients (Ao et al. 2020; Lu et al. 2021). Some classical biomarkers (e.g., CEA, AFP, and CA72-4) have been clinically applied, but their unideal specificity and sensitivity restrict their further utilization (Liu et al. 2022c). Therefore, there is an urgent need in cancer treatment for the exploration of reliable biomarkers with high specificity and sensitivity. FADD has exhibited great potential as a prognostic biomarker for cancer patients due to its aberrant expression pattern in different types of cancer (Table 3).

Chen et al. showed that lung ADC patients with FADD overexpression had a significantly lower overall survival




Fig. 3 Role of FADD in cancer progression. FADD dysregulation contributes to cancer progression by various mechanisms, including crosstalk with oncogenic signaling pathway, induction of inflammation, inhibition of apoptosis, promotion of proliferation, modulation of cell cycle, and enhancement of drug resistance

(OS) rate ($p=0.033$). Multivariate analysis further confirmed that FADD upregulation was an independent poor prognostic biomarker for patients with surgically resected lung ADC ($p=0.027$) (Chen et al. 2021). González-Moles et al. performed a meta-analysis to evaluate the prognostic significance of FADD upregulation in HNSCC. They found that elevated FADD expression was significantly associated with poor OS ($p<0.001$), disease-specific survival (DSS) ($p<0.001$), and disease-free survival (DFS) ($p<0.001$) (Gonzalez-Moles et al. 2020). FADD gene copy number amplification can also be used as a potential prognostic biomarker. For example, Eytan et al. discovered that HNSCC harbored the most frequent genomic amplification of FADD, affecting approximately 30% of patients with poor prognosis (Eytan et al. 2016). Chien et

al. showed that both FADD gene copy number amplification and elevated protein expression were significantly correlated with lymph node metastasis in OSCC patients ($p<0.001$). OSCC patients with both FADD gene copy number amplification and elevated protein expression showed the shortest DFS ($p=0.074$ and $p=0.002$) and OS ($p=0.011$ and $p=0.027$) (Chien et al. 2016). In multiple studies, phosphorylated FADD was identified as a promising prognostic biomarker in distinct types of cancer, including T-cell lymphoblastic lymphoma, PCa, and glottic squamous cell carcinoma (Marin-Rubio et al. 2019a; Matsumura et al. 2009; Schrijvers et al. 2012). In addition, the combination of FADD with other genes can be used to predict cancer treatment prognosis (Cui et al. 2021; Jiang et al. 2021; Yang et al. 2020a). For example, Choi et

Table 3 FADD as a prognostic biomarker in cancer

Cancer types	Clinical values	References
HNSCC	FADD overexpression was significantly associated with worse OS ($p < 0.001$), DSS ($p < 0.001$), and DFS ($p < 0.001$). A 13-gene signature consist of <i>FADD</i> , <i>GABARAPL1</i> , <i>ITGA3</i> , <i>USP10</i> , <i>ST13</i> , <i>MAPK9</i> , <i>PRKN</i> , <i>IKBKB</i> , <i>ITPR1</i> , <i>TP73</i> , <i>MAP2K7</i> , <i>CDKN2A</i> , and <i>EEF2K</i> was an independent prognostic factor ($p < 0.001$) for NSCLC patients. The 13-gene signature significantly stratified HNSCC patients into high- and low-risk groups in terms of OS ($p < 0.0001$). The AUC vale were significant for both the TCGA (0.685) and GEO (0.928). A 3-gene signature consist of <i>FADD</i> , <i>EGFR</i> , and <i>PARK2</i> was an independent prognostic biomarker. ROC analyses revealed high predictive accuracy and sensitivity of the 3-gene signature. A 6-gene signature consist of <i>FADD</i> , <i>EGFR</i> , <i>HSPB8</i> , <i>PRKN</i> , <i>CDKN2A</i> , and <i>ITGA3</i> was an independent prognostic biomarker (AUC=0.709) for NSCLC patients. Lower FADD expression was significantly correlated with better OS ($p=0.0001$) and DFS ($p=0.0006$) in HNSCC patients with lymph node metastasis. A 7-gene signature consist of <i>FADD</i> , <i>ITGA3</i> , <i>CDKN2A</i> , <i>NKX2-3</i> , <i>BAK1</i> , <i>CXCR4</i> , and <i>HSPB8</i> was proved to be effective in predicting the survival rate of HNSCC patients ($p=8.409 \times 10^{-6}$).	Gonzalez-Moles et al. (Gonzalez-Moles et al. 2020) Ren et al. (Ren et al. 2021) Jiang et al. (Jiang et al. 2021) Yang et al. (Yang et al. 2020a) Fan et al. (Fan et al. 2013) Jin et al. (Jin and Qin 2020)
NSCLC	Lung ADC patients with overexpression of FADD exhibited significantly lower OS rates ($p=0.033$). FADD overexpression was an independent poor prognostic biomarker for patients with surgically resected lung ADC ($p=0.027$).	Chen et al. (Chen et al. 2021)
ESCC	A 13-gene signature consist of <i>FADD</i> , <i>PARP1</i> , <i>ITGA6</i> was an independent prognostic factor for ESCC patients. This signature effectively stratified patients in both discovery and validation cohorts by OS ($p=5.162E-8$ and $p=0.052$, respectively).	Cui et al. (Cui et al. 2021)
BC	FADD expression was significantly associated with T stage ($P=0.046$) in BC patients. The combination of <i>FADD</i> , <i>PPFIA1</i> , and <i>TMEM16A</i> gene expressions was significantly correlated with perineural invasion ($p=0.022$). Moreover, the combination of the three gene expressions was closely associated with DFS ($p=0.034$).	Choi et al. (Choi et al. 2014)
Lymphoma	The phosphorylation of FADD (Ser194) was identified as a prognostic biomarker in T-cell lymphoblastic lymphoma. Low or absent expression of the FADD in leukemic cells was identified as a poor independent prognostic biomarker.	Marin-Rubio et al. (Marin-Rubio et al. 2019a) (Tourneur et al. 2004)
Lung cancer	A 6-gene signature consist of <i>FADD</i> , <i>EIF4EBP1</i> , <i>TP63</i> , <i>BNIP3</i> , <i>ATIC</i> , and <i>ERO1A</i> was identified as an independent prognostic biomarker for lung adenocarcinoma and lung squamous cell carcinoma.	Zhu et al. (Zhu et al. 2020)
OSCC	FADD gene copy number and protein expression were identified as potential prognostic biomarkers in patients with OSCC. Patients with both FADD copy number amplification and high protein expression exhibited the shortest DFS ($p=0.074$ and $p=0.002$) and OS ($p=0.011$ and $p=0.027$).	Chien et al. (Chien et al. 2016)

al. demonstrated that FADD expression was significantly associated with the T stage ($p=0.046$) in patients with invasive ductal carcinoma of the breast. Moreover, the combination of *FADD*, *TMEM16A*, and *PPFIA1* expressions showed significant associations with DFS ($P=0.034$) (Choi et al. 2014). Collectively, these studies strongly suggest that FADD possesses great potential as a valuable prognostic biomarker for cancer patients. However, large patient cohorts should be included in future studies to further confirm the application of FADD as a prognostic biomarker in clinics.

FADD as therapeutic target in cancer

FADD has been shown to participate in cancer progression by altering various cellular processes, including proliferation, apoptosis, cell cycle, inflammation, and drug resistance. High FADD expression contributes to the progression of multiple types of cancer, such as PC, OSCC, HNSCC, and BC, indicating that FADD acts as an oncoprotein in these cancers (Chien et al. 2016; Gonzalez-Moles et al. 2020; Zhang et al. 2017; Zhou et al. 2022a).

Thus, a screening or synthesis of novel agents targeting FADD may be an effective therapeutic strategy in these cancers. By contrast, low FADD expression has been observed in several certain types of cancer, such as thymic lymphoma, AML, and GBM, suggesting that FADD serves as a tumor suppressor in these cancers (Newton et al. 2000; Tourneur et al. 2004; Wang et al. 2017). Therefore, the upregulation of FADD in these cancers represents another promising strategy in cancer treatment. In addition, a series of ncRNAs, such as miR-633, H19, and circ 001418, have been identified as upstream regulators of FADD in cancer progression (Pang et al. 2019; Peng et al. 2021; Yan et al. 2017). Targeting these ncRNAs is also an attractive strategy in overcoming cancer. Research on novel agents targeting FADD in cancer treatment remains in the pre-clinical stage. Further studies are required to translate FADD-based therapeutic strategies into clinical utilization in treatment.

Conclusion and perspective

FADD plays a vital role in normal physiological processes by regulating many aspects of cell behavior, such as apoptosis, proliferation, autophagy, and necroptosis. Its expression and activity are tightly controlled through a synergistic regulatory network involving DNA amplification, promoter methylation, gene mutation, TFs, ncRNAs, and PTMs. Therefore, the dysregulation of FADD is closely associated with the initiation and development of many diseases, particularly cancer. FADD dysregulation contributes to cancer progression through various mechanisms, including cross-talking with cancer-related signaling pathways, inhibiting cell apoptosis, facilitating proliferation, disordering cell cycle, inducing inflammation, and enhancing drug resistance. These characteristics endow FADD with great potential as a valuable therapeutic target for a wide range of cancers. Thus, the screening or synthesis of novel agents targeting FADD may be an effective therapeutic strategy for cancer patients. However, current studies are still in the stage of basic and preclinical research. Clinical trials with large patient cohorts should be included as an important direction in future studies. The differential expression of FADD in distinct cancer types also offers an exciting possibility for the development of cancer-tissue-specific treatment plans. For example, FADD is highly expressed in PC, OSCC, BC, and HNSCC, and it promotes the progression of these cancers by serving as an oncoprotein. Targeting oncogenic FADD in these cancer types could be an effective therapeutic strategy for patients. Conversely, FADD is lowly expressed in thymic lymphoma, AML, and GBM, and its overexpression inhibits the progression of these cancers. The upregulation of tumor-suppressive FADD may represent an effective way to overcome these cancer types. In addition, FADD overexpression has been found to enhance drug resistance in a variety of cancers (e.g., GC and PCa), indicating the feasibility of targeting FADD to enhance efficiency in chemotherapy (Pang et al. 2019; Zhang et al. 2017). Accurately evaluating prognoses is a core link in cancer treatment for patients to adjust their therapeutic strategies and prolong their lifespan. The aberrant expression of FADD has been shown to be significantly associated with some pathological features of cancer patients (e.g., OS, DSS and DFS), strongly indicating its potential value as a prognostic biomarker in cancer treatment (Chen et al. 2021; Chien et al. 2016; Gonzalez-Moles et al. 2020). However, there are still some unsolved challenges regarding the strategy of targeting FADD in cancer patients due to the essential role that FADD plays in normal physiological processes. For instance, targeting FADD may cause unexpected abnormal pathological reactions in cancer patients. Nevertheless, FADD has exhibited great potential as a therapeutic target and/or prognostic biomarker in cancer treatment.

Continued in-depth research into its various roles in cancer progression may be of great benefit to the development of FADD-based therapeutic strategies for cancer patients.

Abbreviations

FADD	Fas-associated protein with death domain
DD	death domain
DR	death receptor
NSCLC	non-small cell lung cancer
HCC	hepatocellular carcinoma
DED	death effector domain
TNF	tumor necrosis factor
RIPK	receptor-interacting protein kinase
ATG5	autophagy related-protein 5
NLS	nuclear localization signal
NES	nuclear export signal
DISC	death-inducing signal complex
cFLIP	cellular FLICE (caspase 8)-inhibitory protein
MLKL	mixed lineage kinase domain-like protein
IFN	interferon
Cryo-EM	cryogenic electron microscopy
tDED	tandem DED
TF	transcription factor
ncRNA	non-coding RNAs
PTM	post-translational modifications
MEFs	murine embryonic fibroblasts
BC	breast cancer
BCa	bladder cancer
OSCC	oral squamous cell carcinoma
LUSC	lung squamous cell carcinoma
OC	ovarian cancer
ESCC	esophageal squamous cell carcinoma
HNSCC	head and neck squamous cell carcinoma
CC	cervical cancer
WB	western blotting
FCM	flow cytometry
qRT-PCR	quantitative real-time polymerase chain reaction
CKK-8	cell count kit-8
RT-qPCR	reverse transcription-quantitative polymerase chain reaction
MTT	3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium
Co-IP	co-immunoprecipitation
HIF-1 α	hypoxia-inducible factor-1 α
BRCA1	BC susceptibility gene 1
HRE	hypoxia-responsive element
mRNA	messenger RNA
miRNA	microRNA
lncRNA	long non-coding RNA
circRNA	circular RNA
piRNA	piwi-interacting RNA
3'UTR	3'-untranslated region
GC	gastric cancer
DOX	doxorubicin
CDDP	cisplatin
NP	nucleus pulposus
CK	casein kinase
PLK	polo kinase
AURA	Aurora-A kinase
MKRN1	makorin ring finger protein 1
CHIP	C terminus HSC70-interacting protein
LUBAC	linear ubiquitin chain assembly complex
Drp1	dynamain-related protein 1
GlcNAc	N-acetylglucosamine
AML	acute myeloid leukemia
GBM	glioblastoma
PC	pancreatic cancer
MAPK	mitogen-activated protein kinase
NF- κ B	nuclear factor- κ B
JNK	c-Jun NH2 terminal kinase
ERK	extracellular stress regulated kinase

PTX	paclitaxel
PCa	prostate cancer
MEKK1	MEK kinase 1
TAT	trans-acting activator of transcription
CRC	colorectal cancer
CDK	cyclin dependent kinases
ADC	adenocarcinoma
TMZ	temozolomide
OS	overall survival
DSS	disease-specific survival
DFS	disease-free survival

Acknowledgements

Not applicable.

Author contributions

YL: Writing- Conceptualization, Original draft preparation, Writing - Review & Editing. XL: Data Curation. XZ: Data Curation. JW: Data Curation. XA: Review & Editing, Funding acquisition.

Funding

This work was supported by the China Postdoctoral Science Foundation (2018M642607).

Availability of data and materials

Not applicable.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

All authors have read and approved the final manuscript.

Competing interests

The authors have no conflict of interest.

Received: 12 August 2022 / Accepted: 19 October 2022

Published online: 08 November 2022

References

- Alappat EC, Volkland J, Peter ME. Cell cycle effects by C-FADD depend on its C-terminal phosphorylation site. *J Biol Chem*. 2003;278(43):41585–8.
- Alappat EC, Feig C, Boyerinas B, Volkland J, Samuels M, Murmann AE, et al. Phosphorylation of FADD at serine 194 by CK1 α regulates its nonapoptotic activities. *Mol Cell*. 2005;19(3):321–32.
- Antunovic M, Matic I, Nagy B, Caput Mihalic K, Skelin J, Stambuk J, et al. FADD-deficient mouse embryonic fibroblasts undergo RIPK1-dependent apoptosis and autophagy after NB-UVB irradiation. *J Photochem Photobiol B*. 2019;194:32–45.
- Ao X, Ding W, Ge H, Zhang Y, Ding D, Liu Y. PBX1 is a valuable prognostic biomarker for patients with breast cancer. *Experimental and therapeutic medicine*. 2020;20(1):385–94.
- Bhojani MS, Chen G, Ross BD, Beer DG, Rehemtulla A. Nuclear localized phosphorylated FADD induces cell proliferation and is associated with aggressive lung cancer. *Cell Cycle*. 2005;4(11):1478–81.
- Bowman BM, Sebolt KA, Hoff BA, Boes JL, Daniels DL, Heist KA, et al. Phosphorylation of FADD by the kinase CK1 α promotes KRASG12D-induced lung cancer. *Sci Signal*. 2015;8(361):ra9.
- Brennan K, Lyons C, Fernandes P, Doyle S, Houston A, Brint E. Engagement of Fas differentially regulates the production of LPS-induced proinflammatory cytokines and type I interferons. *FEBS J*. 2019;286(3):523–35.
- Brown LA, Kalloger SE, Miller MA, Shih le M, McKinney SE, Santos JL, et al. Amplification of 11q13 in ovarian carcinoma. *Genes Chromosomes Cancer*. 2008;47(6):481–9.
- Cai F, Xu H, Cao N, Zhang X, Liu J, Lu Y, et al. ADT-OH, a hydrogen sulfide-releasing donor, induces apoptosis and inhibits the development of melanoma in vivo by upregulating FADD. *Cell Death Dis*. 2020;11(1):33.
- Callegari CC, Cavalli JJ, Lima RS, Jucoski TS, Torresan C, Urban CA, et al. Copy number and expression analysis of FOSL1, GSTP1, NTSR1, FADD and CCND1 genes in primary breast tumors with axillary lymph node metastasis. *Cancer Genet*. 2016;209(7–8):331–9.
- Chaitongyot S, Jantaree P, Sokolova O, Naumann M. NF- κ B in Gastric Cancer Development and Therapy. *Biomedicines*. 2021;9(8).
- Chang W, Li M, Song L, Miao S, Yu W, Wang J. Noncoding RNAs from tissue-derived small extracellular vesicles: Roles in diabetes and diabetic complications. *Mol Metab*. 2022a;58:101453.
- Chang W, Wang M, Zhang Y, Yu F, Hu B, Goljanek-Whysall K, et al. Roles of long noncoding RNAs and small extracellular vesicle-long noncoding RNAs in type 2 diabetes. *Traffic*. 2022b.
- Chaudhary PM, Eby MT, Jasmin A, Kumar A, Liu L, Hood L. Activation of the NF- κ B pathway by caspase 8 and its homologs. *Oncogene*. 2000;19(39):4451–60.
- Chen G, Bhojani MS, Heaford AC, Chang DC, Laxman B, Thomas DG, et al. Phosphorylated FADD induces NF- κ B, perturbs cell cycle, and is associated with poor outcome in lung adenocarcinomas. *Proc Natl Acad Sci U S A*. 2005;102(35):12507–12.
- Chen L, Xie G, Feng J, Wen Q, Zang H, Lu J, et al. Overexpression of FADD and Bcl-XS proteins as novel prognostic biomarkers for surgically resected non-small cell lung cancer. *Cancer Biomark*. 2021;30(2):145–54.
- Chien HT, Cheng SD, Chuang WY, Liao CT, Wang HM, Huang SF. Clinical Implications of FADD Gene Amplification and Protein Overexpression in Taiwanese Oral Cavity Squamous Cell Carcinomas. *PLoS ONE*. 2016;11(10):e0164870.
- Choi EJ, Yun JA, Jabeen S, Jeon EK, Won HS, Ko YH, et al. Prognostic significance of TMEM16A, PPF1A1, and FADD expression in invasive ductal carcinoma of the breast. *World J Surg Oncol*. 2014;12:137.
- Choi SG, Kim H, Jeong EI, Lee HJ, Park S, Lee SY, et al. SUMO-Modified FADD Recruits Cytosolic Drp1 and Caspase-10 to Mitochondria for Regulated Necrosis. *Mol Cell Biol*. 2017;37(2).
- Cimino Y, Costes A, Damotte D, Validire P, Mistou S, Cagnard N, et al. FADD protein release mirrors the development and aggressiveness of human non-small cell lung cancer. *Br J Cancer*. 2012;106(12):1989–96.
- Cui H, Weng Y, Ding N, Cheng C, Wang L, Zhou Y, et al. Autophagy-Related Three-Gene Prognostic Signature for Predicting Survival in Esophageal Squamous Cell Carcinoma. *Front Oncol*. 2021;11:650891.
- Debnath S, Sarkar A, Mukherjee DD, Ray S, Mahata B, Mahata T, et al. Eriodictyol mediated selective targeting of the TNFR1/FADD/TRADD axis in cancer cells induce apoptosis and inhibit tumor progression and metastasis. *Transl Oncol*. 2022;21:101433.
- Eberstadt M, Huang B, Chen Z, Meadows RP, Ng SC, Zheng L, et al. NMR structure and mutagenesis of the FADD (Mort1) death-effector domain. *Nature*. 1998;392(6679):941–5.
- Ehlfen H, Krishna-Subramanian S, Ochoa-Callejero L, Kondylis V, Nadi NE, Straub BK, et al. Death receptor-independent FADD signalling triggers hepatitis and hepatocellular carcinoma in mice with liver parenchymal cell-specific NEMO knockout. *Cell Death Differ*. 2014;21(11):1721–32.
- Eytan DF, Snow GE, Carlson S, Derakhshan A, Saleh A, Schiltz S, et al. SMAC Mimetic Birinapant plus Radiation Eradicates Human Head and Neck Cancers with Genomic Amplifications of Cell Death Genes FADD and BIRC2. *Cancer Res*. 2016;76(18):5442–54.
- Fan S, Muller S, Chen ZG, Pan L, Tighiouart M, Shin DM, et al. Prognostic impact of Fas-associated death domain, a key component in death receptor signaling, is dependent on the presence of lymph node metastasis in head and neck squamous cell carcinoma. *Cancer Biol Ther*. 2013;14(4):365–9.
- Fan Y, Li X, Zhang L, Zong Z, Wang F, Huang J, et al. SUMOylation in Viral Replication and Antiviral Defense. *Adv Sci (Weinh)*. 2022;9(7):e2104126.
- Feng S, Ma L, Yang Y, Wu M. Truncated RIP3 (tRIP3) acts upstream of FADD to induce apoptosis in the human hepatocellular carcinoma cell line QGY-7703. *Biochem Biophys Res Commun*. 2006;347(3):558–65.
- Fox JL, Hughes MA, Meng X, Sarnowska NA, Powley IR, Jukes-Jones R, et al. Cryo-EM structural analysis of FADD:Caspase-8 complexes defines the catalytic dimer architecture for co-ordinated control of cell fate. *Nat Commun*. 2021;12(1):819.
- Ge L, Zhang X, Hu S, Song Y, Kong J, Zhang B, et al. H19 suppresses the growth of hepatoblastoma cells by promoting their apoptosis via the signaling pathways of miR-675/FADD and miR-138/PTK2. *J Cell Biochem*. 2019;120(4):5218–31.
- Gibcus JH, Menkema L, Mastik MF, Hermesen MA, de Bock GH, van Velthuisen ML, et al. Amplicon mapping and expression profiling identify the Fas-associated

- death domain gene as a new driver in the 11q13.3 amplicon in laryngeal/pharyngeal cancer. *Clin Cancer Res.* 2007;13(21):6257–66.
- Gomez-Angelats M, Cidlowski JA. Molecular evidence for the nuclear localization of FADD. *Cell Death Differ.* 2003;10(7):791–7.
- Gonzalez-Moles MA, Ayen A, Gonzalez-Ruiz I, de Porrás-Carrique T, Gonzalez-Ruiz L, Ruiz-Avila I, et al. Prognostic and Clinicopathological Significance of FADD Upregulation in Head and Neck Squamous Cell Carcinoma: A Systematic Review and Meta-Analysis. *Cancers.* 2020;12(9).
- Goto E, Tokunaga F. Decreased linear ubiquitination of NEMO and FADD on apoptosis with caspase-mediated cleavage of HOIP. *Biochem Biophys Res Commun.* 2017;485(1):152–9.
- Han B, Meng X, Wu P, Li Z, Li S, Zhang Y, et al. ATRX/EZH2 complex epigenetically regulates FADD/PARP1 axis, contributing to TMZ resistance in glioma. *Theranostics.* 2020;10(7):3351–65.
- Harari-Steinfeld R, Gefen M, Simerzin A, Zorde-Khvaleyevsky E, Rivkin M, Ella E, et al. The lncRNA H19-Derived MicroRNA-675 Promotes Liver Necroptosis by Targeting FADD. *Cancers.* 2021;13(3).
- Hartwig T, Montinaro A, von Karstedt S, Sevko A, Surinova S, Chakravarthy A, et al. The TRAIL-Induced Cancer Secretome Promotes a Tumor-Supportive Immune Microenvironment via CCR2. *Mol Cell.* 2017;65(4):730–42. e5.
- He L, Ren Y, Zheng Q, Wang L, Lai Y, Guan S, et al. Fas-associated protein with death domain (FADD) regulates autophagy through promoting the expression of Ras homolog enriched in brain (Rheb) in human breast adenocarcinoma cells. *Oncotarget.* 2016;7(17):24572–84.
- He X, Peng X, Liu Y, Zhang X, Li H, Yin H. Adenovirus-mediated overexpression FADD induces a significant antitumor effect on human colorectal cancer cells both in vitro and in vivo. *Cell Mol Biol (Noisy-le-grand).* 2018;64(6):31–5.
- Henry CM, Martin SJ. Caspase-8 Acts in a Non-enzymatic Role as a Scaffold for Assembly of a Pro-inflammatory “FADDosome” Complex upon TRAIL Stimulation. *Mol Cell.* 2017;65(4):715–29. e5.
- Hindryckx P, De Vos M, Jacques P, Ferdinande L, Peeters H, Olievier K, et al. Hydroxylase inhibition abrogates TNF-alpha-induced intestinal epithelial damage by hypoxia-inducible factor-1-dependent repression of FADD. *J Immunol.* 2010;185(10):6306–16.
- Hollomon MG, Patterson L, Santiago-O’Farrill J, Kleinerman ES, Gordon N. Knock down of Fas-Associated Protein with Death Domain (FADD) Sensitizes Osteosarcoma to TNFalpha-induced Cell Death. *J Cancer.* 2020;11(7):1657–67.
- Hu WH, Johnson H, Shu HB. Activation of NF-kappaB by FADD, Casper, and caspase-8. *J Biol Chem.* 2000;275(15):10838–44.
- Ikeda T, Tanaka N, Shimada K, Matsumura Y, Miyake M, Anai S, et al. Phosphorylation status of Fas-associated death domain protein is associated with biochemical recurrence after radical prostatectomy. *Urology.* 2013;81(3):607–10.
- Imtiyaz HZ, Zhou X, Zhang H, Chen D, Hu T, Zhang J. The death domain of FADD is essential for embryogenesis, lymphocyte development, and proliferation. *J Biol Chem.* 2009;284(15):9917–26.
- Jang MS, Lee SJ, Kang NS, Kim E. Cooperative phosphorylation of FADD by Aur-A and Plk1 in response to taxol triggers both apoptotic and necrotic cell death. *Cancer Res.* 2011a;71(23):7207–15.
- Jang MS, Lee SJ, Kim CJ, Lee CW, Kim E. Phosphorylation by polo-like kinase 1 induces the tumor-suppressing activity of FADD. *Oncogene.* 2011b;30(4):471–81.
- Jia DD, Jiang H, Zhang YF, Zhang Y, Qian LL, Zhang YF. The regulatory function of piRNA/PIWI complex in cancer and other human diseases: The role of DNA methylation. *Int J Biol Sci.* 2022;18(8):3358–73.
- Jiang Y, Li Y, Ge H, Wu Y, Zhang Y, Guo S, et al. Identification of an autophagy-related prognostic signature in head and neck squamous cell carcinoma. *J Oral Pathol Med.* 2021;50(10):1040–9.
- Jin Y, Qin X. Development of a Prognostic Signature Based on Autophagy-related Genes for Head and Neck Squamous Cell Carcinoma. *Arch Med Res.* 2020;51(8):860–7.
- Katoh M, Katoh M. Comparative genomics on mammalian Fgf3-Fgf4 locus. *Int J Oncol.* 2005;27(1):281–5.
- Kciuk M, Gielecinska A, Budzinska A, Mojzycz M, Kontek R. Metastasis, Pathways MAPK. *International journal of molecular sciences.* 2022;23(7).
- Kondo S, Ishizaka Y, Okada T, Kondo Y, Hitomi M, Tanaka Y, et al. FADD gene therapy for malignant gliomas in vitro and in vivo. *Hum Gene Ther.* 1998;9(11):1599–608.
- Lee EW, Kim JH, Ahn YH, Seo J, Ko A, Jeong M, et al. Ubiquitination and degradation of the FADD adaptor protein regulate death receptor-mediated apoptosis and necroptosis. *Nat Commun.* 2012a;3:978.
- Lee EW, Seo J, Jeong M, Lee S, Song J. The roles of FADD in extrinsic apoptosis and necroptosis. *BMB Rep.* 2012b;45(9):496–508.
- Lin Y, Devin A, Rodriguez Y, Liu ZG. Cleavage of the death domain kinase RIP by caspase-8 prompts TNF-induced apoptosis. *Genes Dev.* 1999;13(19):2514–26.
- Lin Y, Liu J, Chen J, Yao C, Yang Y, Wang J, et al. FADD Phosphorylation Modulates Blood Glucose Levels by Decreasing the Expression of Insulin-Degrading Enzyme. *Mol Cells.* 2020;43(4):373–83.
- Liu Y, Cui H, Huang X, Zhu B, Guan S, Cheng W, et al. MiR-7a is an important mediator in Fas-associated protein with death domain (FADD)-regulated expression of focal adhesion kinase (FAK). *Oncotarget.* 2016;7(32):51393–407.
- Liu Y, Ding W, Ge H, Ponnusamy M, Wang Q, Hao X, et al. FOXK transcription factors: Regulation and critical role in cancer. *Cancer Lett.* 2019;458:1–12.
- Liu Y, Ao X, Ji G, Zhang Y, Yu W, Wang J. Mechanisms of Action And Clinical Implications of MicroRNAs in the Drug Resistance of Gastric Cancer. *Front Oncol.* 2021a;11:768918.
- Liu Y, Ding W, Yu W, Zhang Y, Ao X, Wang J. Long non-coding RNAs: Biogenesis, functions, and clinical significance in gastric cancer. *Mol Ther Oncolytics.* 2021b;23:458–76.
- Liu D, Zhong Z, Karin M. NF-kappaB: A Double-Edged Sword Controlling Inflammation. *Biomedicines.* 2022a;10(6).
- Liu J, Peng Y, Wei W. Cell cycle on the crossroad of tumorigenesis and cancer therapy. *Trends Cell Biol.* 2022b;32(1):30–44.
- Liu Y, Ao X, Jia Y, Li X, Wang Y, Wang J. The FOXO family of transcription factors: key molecular players in gastric cancer. *J Mol Med (Berl).* 2022c;100(7):997–1015.
- Liu Y, Ao X, Wang Y, Li X, Wang J. Long Non-Coding RNA in Gastric Cancer: Mechanisms and Clinical Implications for Drug Resistance. *Front Oncol.* 2022d;12:841411.
- Liu Y, Ao X, Yu W, Zhang Y, Wang J. Biogenesis, functions, and clinical implications of circular RNAs in non-small cell lung cancer. *Mol Ther Nucleic Acids.* 2022e;27:50–72.
- Liu Y, Ao X, Zhou X, Du C, Kuang S. The regulation of PBXs and their emerging role in cancer. *J Cell Mol Med.* 2022f;26(5):1363–79.
- Liu Y, Li Y, Du C, Kuang S, Zhou X, Zhang J, et al. Underlying mechanisms of epithelial splicing regulatory proteins in cancer progression. *J Mol Med (Berl).* 2022g.
- Liu Y, Wang Y, Li X, Jia Y, Wang J, Ao X. FOXO3a in cancer drug resistance. *Cancer Lett.* 2022h;540:215724.
- Lu H, Shi C, Liu X, Liang C, Yang C, Wan X, et al. Identification of ZG16B as a prognostic biomarker in breast cancer. *Open Med (Wars).* 2021;16(1):1–13.
- Luo J, Chen B, Gao CX, Xie HK, Han CN, Zhou CC. SPOP promotes FADD degradation and inhibits NF-kappaB activity in non-small cell lung cancer. *Biochem Biophys Res Commun.* 2018;504(1):289–94.
- Marikar FM, Jin G, Sheng W, Ma D, Hua Z. Metallothionein 2A an interactive protein linking phosphorylated FADD to NF-kappaB pathway leads to colorectal cancer formation. *Chin Clin Oncol.* 2016;5(6):76.
- Marin-Rubio JL, de Arriba MC, Cobos-Fernandez MA, Gonzalez-Sanchez L, Ors I, Sastre I, et al. Deregulated FADD expression and phosphorylation in T-cell lymphoblastic lymphoma. *Oncotarget.* 2016;7(38):61485–99.
- Marin-Rubio JL, Perez-Gomez E, Fernandez-Piqueras J, Villa-Morales M. S194-P-FADD as a marker of aggressiveness and poor prognosis in human T-cell lymphoblastic lymphoma. *Carcinogenesis.* 2019a;40(10):1260–8.
- Marin-Rubio JL, Vela-Martin L, Fernandez-Piqueras J, Villa-Morales M. FADD in Cancer: Mechanisms of Altered Expression and Function, and Clinical Implications. *Cancers.* 2019b;11(10).
- Matsumura Y, Shimada K, Tanaka N, Fujimoto K, Hirao Y, Konishi N. Phosphorylation status of Fas-associated death domain-containing protein regulates telomerase activity and strongly correlates with prostate cancer outcomes. *Pathobiology.* 2009;76(6):293–302.
- Matsuyoshi S, Shimada K, Nakamura M, Ishida E, Konishi N. FADD phosphorylation is critical for cell cycle regulation in breast cancer cells. *Br J Cancer.* 2006;94(4):532–9.
- Meer E, Solanes F, Kohn L, Kuo CY, Wong DA, Pineles S, et al. Ocular findings associated with FADD deficiency resemble familial exudative vitreoretinopathy. *Am J Ophthalmol Case Rep.* 2022;25:101305.
- Mihasan M, Wormwood KL, Sokolowska I, Roy U, Woods AG, Darie CC. Mass Spectrometry- and Computational Structural Biology-Based Investigation of Proteins and Peptides. *Adv Exp Med Biol.* 2019;1140:265–87.
- Milner AE, Palmer DH, Hodgkin EA, Eliopoulos AG, Knox PG, Poole CJ, et al. Induction of apoptosis by chemotherapeutic drugs: the role of FADD in activation of caspase-8 and synergy with death receptor ligands in ovarian carcinoma cells. *Cell Death Differ.* 2002;9(3):287–300.
- Mouasni S, Tourneur L. FADD at the Crossroads between Cancer and Inflammation. *Trends Immunol.* 2018;39(12):1036–53.

- Mrkova Z, Portesova M, Slaninova I. Loss of FADD and Caspases Affects the Response of T-Cell Leukemia Jurkat Cells to Anti-Cancer Drugs. *International journal of molecular sciences*. 2021;22(5).
- Murata M. Inflammation and cancer. *Environ Health Prev Med*. 2018;23(1):50.
- Newton K, Harris AW, Strasser A. FADD/MORT1 regulates the pre-TCR checkpoint and can function as a tumour suppressor. *EMBO J*. 2000;19(5):931–41.
- Nguyen DD, Lee DG, Kim S, Kang K, Rhee S. Integrative Bioinformatics and Functional Analyses of GEO, ENCODE, and TCGA Reveal FADD as a Direct Target of the Tumor Suppressor BRCA1. *International journal of molecular sciences*. 2018;19(5).
- Nussinov R, Tsai CJ, Jang H. Anticancer drug resistance: An update and perspective. *Drug Resist Updat*. 2021;59:100796.
- Osborn SL, Diehl G, Han SJ, Xue L, Kurd N, Hsieh K, et al. Fas-associated death domain (FADD) is a negative regulator of T-cell receptor-mediated necroptosis. *Proc Natl Acad Sci U S A*. 2010;107(29):13034–9.
- Pang X, Zhou Z, Yu Z, Han L, Lin Z, Ao X, et al. Foxo3a-dependent miR-633 regulates chemotherapeutic sensitivity in gastric cancer by targeting Fas-associated death domain. *RNA Biol*. 2019;16(2):233–48.
- Pattje WJ, Melchers LJ, Slagter-Menkema L, Mastik MF, Schrijvers ML, Gibcus JH, et al. FADD expression is associated with regional and distant metastasis in squamous cell carcinoma of the head and neck. *Histopathology*. 2013;63(2):263–70.
- Peng G, Meng H, Pan H, Wang W. CircRNA 001418 Promoted Cell Growth and Metastasis of Bladder Carcinoma via EphA2 by miR-1297. *Curr Mol Pharmacol*. 2021;14(1):68–78.
- Petersen S, Aninat-Meyer M, Schluns K, Gellert K, Dietel M, Petersen I. Chromosomal alterations in the clonal evolution to the metastatic stage of squamous cell carcinomas of the lung. *Br J Cancer*. 2000;82(1):65–73.
- Pyo JO, Jang MH, Kwon YK, Lee HJ, Jun JI, Woo HN, et al. Essential roles of Atg5 and FADD in autophagic cell death: dissection of autophagic cell death into vacuole formation and cell death. *J Biol Chem*. 2005;280(21):20722–9.
- Qin HD, Liao XY, Chen YB, Huang SY, Xue WQ, Li FF, et al. Genomic Characterization of Esophageal Squamous Cell Carcinoma Reveals Critical Genes Underlying Tumorigenesis and Poor Prognosis. *Am J Hum Genet*. 2016;98(4):709–27.
- Ranjan K, Pathak C. Expression of FADD and cFLIPL balances mitochondrial integrity and redox signaling to substantiate apoptotic cell death. *Mol Cell Biochem*. 2016a;422(1–2):135–50.
- Ranjan K, Pathak C. FADD regulates NF-kappaB activation and promotes ubiquitination of cFLIPL to induce apoptosis. *Sci Rep*. 2016b;6:22787.
- Ranjan K, Waghela BN, Vaidya FU, Pathak C. Cell-Penetrable Peptide-Conjugated FADD Induces Apoptosis and Regulates Inflammatory Signaling in Cancer Cells. *International journal of molecular sciences*. 2020;21(18).
- Rasamny JJ, Allak A, Krook KA, Jo VY, Policarpio-Nicolas ML, Sumner HM, et al. Cyclin D1 and FADD as biomarkers in head and neck squamous cell carcinoma. *Otolaryngol Head Neck Surg*. 2012;146(6):923–31.
- Ren Z, Zhang L, Ding W, Luo Y, Shi Z, Shrestha B, et al. Development and validation of a novel survival model for head and neck squamous cell carcinoma based on autophagy-related genes. *Genomics*. 2021;113(1 Pt 2):1166–75.
- Rochat-Steiner V, Becker K, Micheau O, Schneider P, Burns K, Tschopp J. FIST/HIPK3: a Fas/FADD-interacting serine/threonine kinase that induces FADD phosphorylation and inhibits fas-mediated Jun NH(2)-terminal kinase activation. *J Exp Med*. 2000;192(8):1165–74.
- Roy D, Sheng GY, Herve S, Carvalho E, Mahanty A, Yuan S, et al. Interplay between cancer cell cycle and metabolism: Challenges, targets and therapeutic opportunities. *Biomed pharmacotherapy = Biomedecine pharmacotherapie*. 2017;89:288–96.
- Saberi E, Kordi-Tamandani DM, Jamali S, Rigi-Ladiz MA. Analysis of methylation and mRNA expression status of FADD and FAS genes in patients with oral squamous cell carcinoma. *Med Oral Patol Oral Cir Bucal*. 2014;19(6):e562-8.
- Schaub FJ, Han DK, Liles WC, Adams LD, Coats SA, Ramachandran RK, et al. Fas/FADD-mediated activation of a specific program of inflammatory gene expression in vascular smooth muscle cells. *Nat Med*. 2000;6(7):790–6.
- Schrijvers ML, Pattje WJ, Slagter-Menkema L, Mastik MF, Gibcus JH, Langendijk JA, et al. FADD expression as a prognosticator in early-stage glottic squamous cell carcinoma of the larynx treated primarily with radiotherapy. *Int J Radiat Oncol Biol Phys*. 2012;83(4):1220–6.
- Scott NE, Giogha C, Pollock GL, Kennedy CL, Webb AI, Williamson NA, et al. The bacterial arginine glycosyltransferase effector NleB preferentially modifies Fas-associated death domain protein (FADD). *J Biol Chem*. 2017;292(42):17337–50.
- Seo J, Lee EW, Shin J, Seong D, Nam YW, Jeong M, et al. K6 linked polyubiquitylation of FADD by CHIP prevents death inducing signaling complex formation suppressing cell death. *Oncogene*. 2018;37(36):4994–5006.
- Sheikh MS, Huang Y. The FADD is going nuclear. *Cell Cycle*. 2003;2(4):346–7.
- Shimada K, Matsuyoshi S, Nakamura M, Ishida E, Kishi M, Konishi N. Phosphorylation of FADD is critical for sensitivity to anticancer drug-induced apoptosis. *Carcinogenesis*. 2004;25(7):1089–97.
- Shimada K, Matsuyoshi S, Nakamura M, Ishida E, Konishi N. Phosphorylation status of Fas-associated death domain-containing protein (FADD) is associated with prostate cancer progression. *J Pathol*. 2005;206(4):423–32.
- Simmons CPL, McMillan DC, McWilliams K, Sande TA, Fearon KC, Tuck S, et al. Prognostic Tools in Patients With Advanced Cancer: A Systematic Review. *J Pain Symptom Manage*. 2017;53(5):962–70. e10.
- Singh N, Baby D, Rajguru JP, Patil PB, Thakkannavar SS, Pujari VB. Inflammation and cancer. *Ann Afr Med*. 2019;18(3):121–6.
- Singh P, Lim B. Targeting Apoptosis in Cancer. *Curr Oncol Rep*. 2022;24(3):273–84.
- Soung YH, Lee JW, Kim SY, Nam SW, Park WS, Kim SH, et al. Mutation of FADD gene is rare in human colon and stomach cancers. *APMIS*. 2004;112(9):595–7.
- Sprick MR, Weigand MA, Rieser E, Rauch CT, Juo P, Blenis J, et al. FADD/MORT1 and caspase-8 are recruited to TRAIL receptors 1 and 2 and are essential for apoptosis mediated by TRAIL receptor 2. *Immunity*. 2000;12(6):599–609.
- Srivatsan ES, Chakrabarti R, Zainabadi K, Pack SD, Benyamini P, Mendonca MS, et al. Localization of deletion to a 300 Kb interval of chromosome 11q13 in cervical cancer. *Oncogene*. 2002;21(36):5631–42.
- Sun BH, Zhao XP, Wang BJ, Yang DL, Hao LJ. FADD and TRADD expression and apoptosis in primary hepatocellular carcinoma. *World J Gastroenterol*. 2000;6(2):223–7.
- Sung TC, Chen Z, Thuret S, Vilar M, Gage FH, Riek R, et al. P45 forms a complex with FADD and promotes neuronal cell survival following spinal cord injury. *PLoS ONE*. 2013;8(7):e69286.
- Tewari R, Sharma V, Koul N, Sen E. Involvement of mitofosin-mediated ERK activation in glioma cell apoptosis through Fas regulation. *J Neurochem*. 2008;107(3):616–27.
- Tourneur L, Delluc S, Levy V, Valensi F, Radford-Weiss I, Legrand O, et al. Absence or low expression of fas-associated protein with death domain in acute myeloid leukemia cells predicts resistance to chemotherapy and poor outcome. *Cancer Res*. 2004;64(21):8101–8.
- Tummers B, Mari L, Guy CS, Heckmann BL, Rodriguez DA, Ruhl S, et al. Caspase-8-Dependent Inflammatory Responses Are Controlled by Its Adaptor, FADD, and Necroptosis. *Immunity*. 2020;52(6):994–1006. e8.
- Vanden Berghe T, van Loo G, Saelens X, Van Gurp M, Brouckaert G, Kalai M, et al. Differential signaling to apoptotic and necrotic cell death by Fas-associated death domain protein FADD. *J Biol Chem*. 2004;279(9):7925–33.
- Verboom L, Martens A, Priem D, Hoste E, Sze M, Vikkula H, et al. OTULIN Prevents Liver Inflammation and Hepatocellular Carcinoma by Inhibiting FADD- and RIPK1 Kinase-Mediated Hepatocyte Apoptosis. *Cell Rep*. 2020;30(7):2237–47. e6.
- Vesely ED, Heilig CW, Brosius FC. 3rd. GLUT1-induced cFLIP expression promotes proliferation and prevents apoptosis in vascular smooth muscle cells. *Am J Physiol Cell Physiol*. 2009;297(3):C759-65.
- Villa-Morales M, Gonzalez-Gugel E, Shahbazi MN, Santos J, Fernandez-Piqueras J. Modulation of the Fas-apoptosis-signalling pathway by functional polymorphisms at Fas, FasL and Fadd and their implication in T-cell lymphoblastic lymphoma susceptibility. *Carcinogenesis*. 2010;31(12):2165–71.
- Vilmont V, Filhol O, Hesse AM, Coute Y, Hue C, Remy-Tourneur L, et al. Modulatory role of the anti-apoptotic protein kinase CK2 in the sub-cellular localization of Fas associated death domain protein (FADD). *Biochim Biophys Acta*. 2015;1853(11 Pt A):2885–96.
- Wang HQ, Yu XD, Liu ZH, Cheng X, Samartzis D, Jia LT, et al. Deregulated miR-155 promotes Fas-mediated apoptosis in human intervertebral disc degeneration by targeting FADD and caspase-3. *J Pathol*. 2011a;225(2):232–42.
- Wang QR, Yao XQ, Wen G, Fan Q, Li YJ, Fu XQ, et al. Apigenin suppresses the growth of colorectal cancer xenografts via phosphorylation and up-regulated FADD expression. *Oncol Lett*. 2011b;2(1):43–7.
- Wang JX, Zhang XJ, Li Q, Wang K, Wang Y, Jiao JQ, et al. MicroRNA-103/107 Regulate Programmed Necrosis and Myocardial Ischemia/Reperfusion Injury Through Targeting FADD. *Circul Res*. 2015;117(4):352–63.
- Wang HB, Li T, Ma DZ, Ji YX, Zhi H. Overexpression of FADD and Caspase-8 inhibits proliferation and promotes apoptosis of human glioblastoma cells. *Biomed pharmacotherapy = Biomedecine pharmacotherapie*. 2017;93:1–7.
- Wang M, Yu F, Li P, Wang K. Emerging Function and Clinical Significance of Exosomal circRNAs in Cancer. *Mol Ther Nucleic Acids*. 2020;21:367–83.

- Wang M, Chen X, Yu F, Ding H, Zhang Y, Wang K. Extrachromosomal Circular DNAs: Origin, formation and emerging function in Cancer. *Int J Biol Sci.* 2021;17(4):1010–25.
- Wang M, Yu F, Zhang Y, Chang W, Zhou M. The Effects and Mechanisms of Flavonoids on Cancer Prevention and Therapy: Focus on Gut Microbiota. *Int J Biol Sci.* 2022;18(4):1451–75.
- Wen ZJ, Xin H, Wang YC, Liu HW, Gao YY, Zhang YF. Emerging roles of circRNAs in the pathological process of myocardial infarction. *Mol Ther Nucleic Acids.* 2021;26:828–48.
- Wichnieski C, Maheshwari K, Souza LC, Nieves F, Tartari T, Garlet GP, et al. DNA methylation profiles of immune response-related genes in apical periodontitis. *Int Endod J.* 2019;52(1):5–12.
- Xia C, Dong X, Li H, Cao M, Sun D, He S, et al. Cancer statistics in China and United States, 2022: profiles, trends, and determinants. *Chin Med J (Engl).* 2022;135(5):584–90.
- Xu H, He L, Feng X, Kapoor A, Tang D. Specific reduction of fas-associated protein with death domain (FADD) in clear cell renal cell carcinoma. *Cancer Invest.* 2009;27(8):836–43.
- Yamada N, Noguchi S, Kumazaki M, Shinohara H, Miki K, Naoe T, et al. Epigenetic regulation of microRNA-128a expression contributes to the apoptosis-resistance of human T-cell leukaemia jurkat cells by modulating expression of fas-associated protein with death domain (FADD). *Biochim Biophys Acta.* 2014;1843(3):590–602.
- Yan J, Zhang Y, She Q, Li X, Peng L, Wang X, et al. Long Noncoding RNA H19/miR-675 Axis Promotes Gastric Cancer via FADD/Caspase 8/Caspase 3 Signaling Pathway. *Cell Physiol Biochem.* 2017;42(6):2364–76.
- Yang YW, Zhang CM, Huang XJ, Zhang XX, Zhang LK, Li JH, et al. Tumor-targeted delivery of a C-terminally truncated FADD (N-FADD) significantly suppresses the B16F10 melanoma via enhancing apoptosis. *Sci Rep.* 2016;6:34178.
- Yang C, Mei H, Peng L, Jiang F, Xie B, Li J. Prognostic Correlation of an Autophagy-Related Gene Signature in Patients with Head and Neck Squamous Cell Carcinoma. *Comput Math Methods Med.* 2020a;2020:7397132.
- Yang C, Ran Q, Zhou Y, Liu S, Zhao C, Yu X, et al. Doxorubicin sensitizes cancer cells to Smac mimetic via synergistic activation of the CYLD/RIPK1/FADD/caspase-8-dependent apoptosis. *Apoptosis.* 2020b;25(5–6):441–55.
- Yin A, Jiang Y, Zhang X, Luo H. Overexpression of FADD enhances 5-fluorouracil-induced apoptosis in colorectal adenocarcinoma cells. *Med Oncol.* 2010;27(2):397–405.
- Yoo NJ, Lee SH, Jeong EG, Lee JW, Soung YH, Nam SW, et al. Expression of nuclear and cytoplasmic phosphorylated FADD in gastric cancers. *Pathol Res Pract.* 2007;203(2):73–8.
- Yuan XJ, Whang YE. PTEN sensitizes prostate cancer cells to death receptor-mediated and drug-induced apoptosis through a FADD-dependent pathway. *Oncogene.* 2002;21(2):319–27.
- Zaafan MA, Abdelhamid AM. The cardioprotective effect of microRNA-103 inhibitor against isoprenaline-induced myocardial infarction in mice through targeting FADD/RIPK pathway. *Eur Rev Med Pharmacol Sci.* 2021;25(2):837–44.
- Zhang X, Vallabhaneni R, Loughran PA, Shapiro R, Yin XM, Yuan Y, et al. Changes in FADD levels, distribution, and phosphorylation in TNFalpha-induced apoptosis in hepatocytes is caspase-3, caspase-8 and BID dependent. *Apoptosis.* 2008;13(8):983–92.
- Zhang J, Pan C, Xu T, Niu Z, Ma C, Xu C. Interleukin 18 augments growth ability via NF-kappaB and p38/ATF2 pathways by targeting cyclin B1, cyclin B2, cyclin A2, and Bcl-2 in BRL-3A rat liver cells. *Gene.* 2015;563(1):45–51.
- Zhang R, Liu Y, Hammache K, He L, Zhu B, Cheng W, et al. The role of FADD in pancreatic cancer cell proliferation and drug resistance. *Oncol Lett.* 2017;13(3):1899–904.
- Zhang L, Zhang Y, Zhao Y, Wang Y, Ding H, Xue S, et al. Circulating miRNAs as biomarkers for early diagnosis of coronary artery disease. *Expert Opin Ther Pat.* 2018;28(8):591–601.
- Zhang L, Zhang Y, Wang Y, Zhao Y, Ding H, Li P. Circular RNAs: Functions and Clinical Significance in Cardiovascular Disease. *Front Cell Dev Biol.* 2020a;8:584051.
- Zhang L, Zhang Y, Xue S, Ding H, Wang Y, Qi H, et al. Clinical significance of circulating microRNAs as diagnostic biomarkers for coronary artery disease. *J Cell Mol Med.* 2020b;24(1):1146–50.
- Zhang YF, Shan C, Wang Y, Qian LL, Jia DD, Zhang YF, et al. Cardiovascular toxicity and mechanism of bisphenol A and emerging risk of bisphenol S. *The Science of the total environment.* 2020c;723:137952.
- Zhang L, Wang Y, Yu F, Li X, Gao H, Li P. CircHIPK3 Plays Vital Roles in Cardiovascular Disease. *Front Cardiovasc Med.* 2021a;8:733248.
- Zhang Y, Jia DD, Zhang YF, Cheng MD, Zhu WX, Li PF, et al. The emerging function and clinical significance of circRNAs in Thyroid Cancer and Autoimmune Thyroid Diseases. *Int J Biol Sci.* 2021b;17(7):1731–41.
- Zhang Y, Wei YJ, Zhang YF, Liu HW, Zhang YF. Emerging Functions and Clinical Applications of Exosomal ncRNAs in Ovarian Cancer. *Front Oncol.* 2021c;11:765458.
- Zhang C, Liu X, Jin S, Chen Y, Guo R. Ferroptosis in cancer therapy: a novel approach to reversing drug resistance. *Mol Cancer.* 2022;21(1):47.
- Zhao X, Yin H, Li N, Zhu Y, Shen W, Qian S, et al. An Integrated Regulatory Network Based on Comprehensive Analysis of mRNA Expression, Gene Methylation and Expression of Long Non-coding RNAs (lncRNAs) in Myelodysplastic Syndromes. *Front Oncol.* 2019;9:200.
- Zhou R, Zhu X, Peng Y, Zhong L, Peng L, Yang B, et al. Clinical Impact of 11q13.3 Amplification on Immune Cell Infiltration and Prognosis in Breast Cancer. *Int J Gen Med.* 2022a;15:4037–52.
- Zhou W, Lai Y, Zhu J, Xu X, Yu W, Du Z, et al. The Classical Apoptotic Adaptor FADD Regulates Glycolytic Capacity in Acute Lymphoblastic Leukemia. *Int J Biol Sci.* 2022b;18(8):3137–55.
- Zhou X, Ao X, Jia Z, Li Y, Kuang S, Du C, et al. Non-coding RNA in cancer drug resistance: Underlying mechanisms and clinical applications. *Front Oncol.* 2022c;12:951864.
- Zhu GF, Yang LX, Guo RW, Liu H, Shi YK, Wang H, et al. miR-155 inhibits oxidized low-density lipoprotein-induced apoptosis of RAW264.7 cells. *Mol Cell Biochem.* 2013;382(1–2):253–61.
- Zhu J, Wang M, Hu D. Development of an autophagy-related gene prognostic signature in lung adenocarcinoma and lung squamous cell carcinoma. *PeerJ.* 2020;8:e8288.
- Zuo YB, Zhang YF, Zhang R, Tian JW, Lv XB, Li R, et al. Ferroptosis in Cancer Progression: Role of Noncoding RNAs. *Int J Biol Sci.* 2022;18(5):1829–43.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.